antibodies .- online.com





Datasheet for ABIN7520036

Galectin 3 Protein (LGALS3)



Overview

Quantity:	20 μg
Target:	Galectin 3 (LGALS3)
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Biological Activity:	Active

Product Details

Product Details		
Purpose:	Active Recombinant Human Galectin-3/LGALS3 Protein	
Sequence:	ADNFSLHDAL SGSGNPNPQG WPGAWGNQPA GAGGYPGASY PGAYPGQAPP GAYPGQAPPG AYPGAPGAYP GAPAPGVYPG PPSGPGAYPS SGQPSATGAY PATGPYGAPA GPLIVPYNLP LPGGVVPRML ITILGTVKPN ANRIALDFQR GNDVAFHFNP RFNENNRRVI VCNTKLDNNW	
	GREERQSVFP FESGKPFKIQ VLVEPDHFKV AVNDAHLLQY NHRVKKLNEI SKLGISGDID LTSASYTMI	
Specificity:	Ala2-Ile250	
Purity:	> 97 % by SDS-PAGE.	
Sterility:	0.22 µm filtered	
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.	
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized Human Galectin-3 at 1 μg/ml (100 μL/well) can bind Galectin-3/LGALS3 Rabbit mAb with a linear range of 0.24-35 ng/mL.	

Target Details

rarget betails	
Target:	Galectin 3 (LGALS3)
Alternative Name:	Galectin-3/LGALS3 (LGALS3 Products)
Target Type:	Chemical
Background:	Description: This protein is a member of the galectin family of carbohydrate binding proteins. Members of this protein family have an affinity for beta-galactosides. The encoded protein is characterized by an N-terminal proline-rich tandem repeat domain and a single C-terminal carbohydrate recognition domain. This protein can self-associate through the N-terminal domain allowing it to bind to multivalent saccharide ligands. This protein localizes to the extracellular matrix, the cytoplasm and the nucleus. This protein plays a role in numerous cellular functions including apoptosis, innate immunity, cell adhesion and T-cell regulation. The protein exhibits antimicrobial activity against bacteria and fungi. Name: CBP35, GAL3, GALBP, GALIG, L31, LGALS2, MAC2, LGALS3, GAL3, GALBP, GALIG, L31, LGALS2, MAC2, LGALS3, GAL3, GALBP, GALIG, L31, LGALS2, MAC2, LGALS3, GAL3, GALBP, GALIG, L31, LGALS2, MAC2
Gene ID:	3958
UniProt:	P17931
Pathways:	RTK Signaling
Application Details	
Restrictions:	For Research Use only
Handling	

Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.