antibodies -online.com





Datasheet for ABIN7520111 **IFNA1 Protein (His tag)**

Go to Product page

Overview

| Quantity: | 50 μg |
|-------------------------------|--|
| Target: | IFNA1 |
| Origin: | Mouse |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This IFNA1 protein is labelled with His tag. |
| | |

Product Details

| Purpose: | Recombinant Mouse IFNA1 Protein |
|------------------|---|
| Sequence: | CDLPQTHNLR NKRALTLLVQ MRRLSPLSCL KDRKDFGFPQ EKVDAQQIKK AQAIPVLSEL TQQILNIFTS KDSSAAWNTT LLDSFCNDLH QQLNDLQGCL MQQVGVQEFP LTQEDALLAV |
| | RKYFHRITVY LREKKHSPCA WEVVRAEVWR ALSSSANVLG RLREEK |
| Specificity: | Cys24-Lys189 |
| Purity: | > 97 % by SDS-PAGE. |
| Sterility: | 0.22 μm filtered |
| Endotoxin Level: | < 0.1EU/µg |

Target Details

| Target: | IFNA1 |
|-------------------|------------------------|
| Alternative Name: | IFNA1 (IFNA1 Products) |

Target Details

Description: IFNA1, also known as IFN-alpha and IFNA, belongs to the alpha/beta interferon family. Interferons(IFNs) are proteins made and released by host cells in response to the presence of pathogens such as viruses, bacteria, parasites, or tumor cells. They belong to the large class of glycoproteins known as cytokines. IFNs stimulate the production of two enzymes: a protein kinase and an oligoadenylate synthetase. They allow for communication between cells to trigger the protective defenses of the immune system that eradicate pathogens or tumors. IFNs can activate immune cells, such as natural killer cells and macrophages, they increase recognition of infection or tumor cells by up-regulating antigen presentation to T lymphocytes, and they also increase the ability of uninfected host cells to resist new infection by the virus. Leukocyte interferon is produced predominantly by B lymphocytes. Immune interferon is produced by mitogen- or antigen-stimulated T lymphocytes. IFNA1 is produced by macrophages and has antiviral activities.

Name: Ifa1, If1ai14,IFNA1

Gene ID:

15962

UniProt:

P01572

Pathways:

JAK-STAT Signaling, Hepatitis C

Application Details

Restrictions:

For Research Use only

Handling

| Format: | Lyophilized |
|------------------|---|
| Reconstitution: | Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles. |
| Concentration: | 1.65 mg/mL |
| Buffer: | Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. |