

Datasheet for ABIN7520120

FIL1d Protein



Overview

Quantity:	10 μg
Target:	FIL1d
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant

Product Details

Purpose:	Recombinant Mouse IL-36Ra/IL-1F5 Protein
Sequence:	VLSGALCFRM KDSALKVLYL HNNQLLAGGL HAEKVIKGEE ISVVPNRALD ASLSPVILGV QGGSQCLSCG TEKGPILKLE PVNIMELYLG AKESKSFTFY RRDMGLTSSF ESAAYPGWFL CTSPEADQPV RLTQIPEDPA WDAPITDFYF QQCD
Specificity:	Val3-Asp156
Purity:	> 97 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	601EU/µg

Target Details

Target:	FIL1d
Alternative Name:	IL-36Ra/IL-1F5 (FIL1d Products)
Background:	Description: Interleukin-1 family member 5 (IL-1F5), also known as interleukin 36 receptor
	antagonist (IL36RA), is a member of the interleukin 1 cytokine family. This cytokine was shown

to specifically inhibit the activation of NF-kappaB induced by interleukin 1 family, member 6 (IL1F6). IL-1F5 is a highly and specific antagonist of the IL-1 receptor-related protein 2-mediated response to interleukin 1 family member 9 (IL1F9). IL-1F5 could constitute part of an independent signaling system analogous to interleukin-1 alpha (IL-1A), beta (IL-1B) receptor agonist, and interleukin-1 receptor type I (IL-1R1), which is present in epithelial barriers and takes part in the local inflammatory response. It has been proved that IL-1F5 induces IL-4 mRNA and protein expression in glia in vitro and enhances hippocampal expression of IL-4 following intracerebroventricular injection. The inhibitory effect of IL-1F5 on LPS-induced IL-1β is attenuated in cells from IL-4-defective mice. Experiment results suggest that IL-1F5 mediates anti-inflammatory effects through its ability to induce IL-4 production and that this is a consequence of its interaction with the orphan receptor, single Ig IL-1R-related molecule (SIGIRR)/TIR8, as the effects were not observed in SIGIRR-/- mice. In contrast to its effects in brain tissue, IL-1F5 did not attenuate LPS-induced changes, or up-regulated IL-4 in macrophages or dendritic cells, suggesting that the effect is confined to the brain.

Gene ID:

54450

UniProt:

Q9QYY1

Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile
	distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is
	recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 %
	Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.