

Datasheet for ABIN7520184 **IL2RG Protein (Fc Tag,AVI tag)**

[Go to Product page](#)

Overview

Quantity:	10 µg
Target:	IL2RG
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL2RG protein is labelled with Fc Tag,AVI tag.

Product Details

Purpose:	Recombinant Cynomolgus IL-2RG/CD132 Protein
Sequence:	LNTTILTPNG NEDATDFFL TSMPTDSLVS STLPLPEVQC FVFNVEYMNC TWNSSSEPQP TNLTLHYWYK NSDNDKVQKC SHYLFSEEIT SGCQLQEKEI HLYQTFVVL QDPREPRRQA TQMLKLQNLV IPWAPENLTL RKLSESQLEL NWNNRFLNHC LEHLVQYRTD WDHWSWTEQSV DYRHKFSLPS VDGQKRYTFR VRSRFNPLCG SAQHWSEWSH PIHWGSNSSK EN
Specificity:	Leu23-Asn254
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.

Target Details

Target:	IL2RG
Alternative Name:	IL-2RG/CD132 (IL2RG Products)
Background:	Description: The common gamma chain (γc) (or CD132), also known as interleukin-2 receptor

Target Details

subunit gamma or IL2RG, is a member of the type I cytokine receptor family expressed on most lymphocyte (white blood cell) populations, and its gene is found on the X-chromosome of mammals. The common gamma chain (γ c) (or IL2RG), is a cytokine receptor subunit that is common to the receptor complexes for at least six different interleukin receptors: IL-2, IL-4, IL-7, IL-9, IL-15, and the interleukin-21 receptor. It is a component of multiple cytokine receptors that are essential for lymphocyte development and function. X-linked severe combined immunodeficiency (X-SCID) is a rare and potentially fatal disease caused by mutations of IL2RG, the gene encoding IL2RG. IL2RG was demonstrated to be a component of the IL-4 receptor based on chemical cross-linking data, the ability of IL2RG to augment IL-4 binding affinity. The observation that IL-2R gamma is a functional component of the IL-4 receptor, together with the finding that IL-2R gamma associates with the IL-7 receptor, begins to elucidate why a deficiency of this common gamma chain (gamma c) has a profound effect on lymphoid function and development, as seen in X-linked severe combined immunodeficiency.

Name: IL2RG,CD132,CIDX,IMD4,P64,SCIDX,SCIDX1,gammaC,IL2RG

Gene ID: 641338

UniProt: [Q38JL2](#)

Pathways: [JAK-STAT Signaling, Growth Factor Binding](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Buffer: Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4.

Storage: -20 °C, -80 °C

Storage Comment: Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.