antibodies -online.com





IL-9 Protein (His tag)



Overview

| Quantity: | 20 μg |
|-------------------------------|---|
| Target: | IL-9 (IL9) |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Biological Activity: | Active |
| Purification tag / Conjugate: | This IL-9 protein is labelled with His tag. |

Product Details

| Purpose: | Active Recombinant Human IL-9 Protein |
|------------------------------|--|
| Sequence: | QGCPTLAGIL DINFLINKMQ EDPASKCHCS ANVTSCLCLG IPSDNCTRPC FSERLSQMTN TTMQTRYPLI FSRVKKSVEV LKNNKCPYFS CEQPCNQTTA GNALTFLKSL LEIFQKEKMR GMRGKI |
| Specificity: | Gln19-Ile144 |
| Purity: | > 97 % by SDS-PAGE. |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | <0.1EU/µg |
| Biological Activity Comment: | Measured in a cell proliferation assay using M07e human megakaryocytic leukemic cells. Avanzi, G. et al. (1988) Br. J. Haematol. 69:359. The ED ₅₀ for this effect is 0.48-1.91 ng/mL. |

Buffer:

Storage:

| Target Details | |
|---------------------|---|
| Target: | IL-9 (IL9) |
| Alternative Name: | IL-9 (IL9 Products) |
| Background: | Description: Interleukin 9, also known as IL-9, is a cytokine (cell signaling molecule) belonging to the group of interleukins. IL-9 is a cytokine that acts as a regulator of a variety of hematopoietic cells. This cytokine stimulates cell proliferation and prevents apoptosis. It functions through the interleukin 9 receptor (IL-9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. Name: P40, HP40, IL-9,IL9 |
| Gene ID: | 3578 |
| UniProt: | P15248 |
| Pathways: | JAK-STAT Signaling |
| Application Details | |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Reconstitution: | Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % |

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

-20 °C,-80 °C

Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.