

[Go to Product page](#)

Datasheet for ABIN7520333 MAPT Protein (His tag)

Overview

Quantity:	10 µg
Target:	MAPT
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MAPT protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human Microtubule-associated protein tau/MAPT Protein
Sequence:	MAEPRQEFEV MEDHAGTYGL GDRKDQGGYT MHQDQEGDTD AGLKESPLQT PTEDGSEEPG SETSDAKSTP TAEDVTAPLV DEGAPGKQAA AQPHTIEPEG TTAEAEAGIGD TPSLEDEAAG HVTQARMVSK SKDGTGSDDK KAKGADGKTK IATPRGAAPP GQKGQANATR IPAKTTPAPK TPPSSGEPPK SGDRSGYSSP GSPGTPGSRS RTPSLPTPPT REPKKVAVVR TPPKSPSSAK SRLQTAPVPM PDLKNVSKI GSTENLKHQP GGGKVQIINK KLDLSNVQSK CGSKDNIKHV PGGGSVQIVY KPDLSKVTS KCGSLGNIHH KPGGGQVEVK SEKLDFKDRV QSKIGSLDNI THVPGGGNKK IETHKLTFRE NAKAKTDHGA EIVYKSPVVS GDTSPRHLSN VSSTGSIDMV DSPQLATLAD EVSASLAKQG L
Specificity:	Met1-Leu441
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	<0.1EU/µg

Target Details

Target:	MAPT
Alternative Name:	Microtubule-associated protein tau/MAPT (MAPT Products)
Background:	<p>Description: MAPT (microtubule-associated protein tau) can produce tau proteins. Tau proteins are proteins that stabilize microtubules. They are abundant in neurons of the central nervous system and are less common elsewhere, but are also expressed at very low levels in CNS astrocytes and oligodendrocytes. When tau proteins are defective, and no longer stabilize microtubules properly, they can result in dementias such as Alzheimer"s disease. Tau protein is a highly soluble microtubule-associated protein (MAP). In humans, these proteins are mostly found in neurons compared to non-neuronal cells. One of tau"s main functions is to modulate the stability of axonal microtubules. Other nervous system MAPs may perform similar functions, as suggested by tau knockout mice, who did not show abnormalities in brain development - possibly because of compensation in tau deficiency by other MAPs. Tau-F is known as "2N4R," "Isoform Tau-F," "Tau-4" or "Tau 441", which consisting of 441 amino acid. Tau-F is a potential therapeutic target for pathogenesis.</p> <p>Name: DDPAC, FTDP-17, MAPTL, MSTD, MTBT1, MTBT2, PPND, PPP1R103, TAU,MAPT,FTDP-17,MAPTL,MSTD,MTBT1,MTBT2,PPND,PPP1R103,TAU,Tau, DDPAC, microtubule-associated protein tau</p>
Gene ID:	4137
UniProt:	P10636-8
Pathways:	MAPK Signaling , Microtubule Dynamics , M Phase , Regulation of Cell Size

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C

Handling

Storage Comment: Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.