# antibodies -online.com





## Datasheet for ABIN7520344

## NCR1 Protein (rFc Tag)



#### Overview

Quantity:	10 μg
Target:	NCR1
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NCR1 protein is labelled with rFc Tag.

### **Product Details**

Purpose:	Recombinant Mouse NCR1/CD335 Protein
Sequence:	QRINTEKETL PKPIIWAKPS IMVTNGNSVN IWCQGAQSAS EYQLYFEGSF FALERPKPSR
	SMNKVRFFIS QMTSHTAGIY TCFYQSGELW SKSSNPLKLV VTGLYDTPNL WVYPRPEVTL
	GENVTFFCQL KTATSKFFLL KERGSNHIQN KYGNIQAEFP MGPVTRAHRG TYRCFGSYND
	YAWSFPSEPV TLLITGGVEN SSLAPTDPTS SLDYWEFDLS TNESGLQKDS AFWDHTTQN
Specificity:	Gln17-Asn255
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	<0.1EU/µg

## **Target Details**

Target:	NCR1
Alternative Name:	NCR1/CD335 (NCR1 Products)

## **Target Details**

Background:	Description: NCR1, also known as NK-p46 and CD335, is a natural cytotoxicity receptor(NCR).
	NCRs are type I transmembrane proteins with 1-2 extracellular immunoglobulin domains, a
	transmembrane domain containing a positively charged amino acid residue, and a short
	cytoplasmic tail. All are expressed almost exclusively by NK cells and play a major role in
	triggering NK-mediated killing of most tumor cell lines. NKp46 has two extracellular Ig-like
	domains followed by a $\sim\!40$ residue stalk region, a type I transmembrane domain, and a short
	cytoplasmic tail. NKp46 has been implicated in NK cell-mediated lysis of several autologous
	tumor cells, pathogen-infected cell lines, and mononuclear phagocytes infected with an
	intracellular bacterium.
	Name: Ly94,NKp46,Ncr1
Gene ID:	17086
UniProt:	Q8C567
Pathways:	Regulation of Leukocyte Mediated Immunity
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile
	distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is
	recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 %
	Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein
	solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.