

[Go to Product page](#)

## Datasheet for ABIN7520383 PTH1R Protein (His tag)

### Overview

Quantity:	20 µg
Target:	PTH1R
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PTH1R protein is labelled with His tag.

### Product Details

Purpose:	Recombinant Human PTH1R Protein
Sequence:	DADDVMTKEE QIFLLHRAQA QCEKRLKEVL QRPASIMESD KGWTSASTSG KPRKDKASGK LYPESEEDKE APTGSRYSRGR PCLPEWDHIL CWPLGAPGEV VAVPCPDYIY DFNHKGHAYR RCDRNGSWEL VPGHNRTWAN YSECVKFLTN ETREREVFDR LG
Specificity:	Asp27-Gly188
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	<0.1EU/µg

### Target Details

Target:	PTH1R
Alternative Name:	PTH1R ( <a href="#">PTH1R Products</a> )

## Target Details

Background:	<p>Description: Parathyroid hormone / parathyroid hormone-related peptide receptor, also known as PTH / PTHrP type I receptor, PTH/PTHr receptor, Parathyroid hormone 1 receptor, PTH1 receptor, PTH1R and PTHR, is a multi-pass membrane protein which belongs to the G-protein coupled receptor 2 family. PTH1R is expressed in most tissues. It is most abundant in kidney, bone and liver. PTH1R is expressed in high levels in bone and kidney and regulates calcium ion homeostasis through activation of adenylate cyclase and phospholipase C. In bone, PTH1R is expressed on the surface of osteoblasts. When the receptor is activated, these cells in turn stimulate osteoclasts to ultimately increase the resorption rate. PTH1R is a receptor for parathyroid hormone and for parathyroid hormone-related peptide. The activity of PTH1R is mediated by G proteins which activate adenylyl cyclase and also a phosphatidylinositol-calcium second messenger system. Defects in PTH1R are the cause of Jansen metaphyseal chondrodysplasia (JMC), chondrodysplasia Blomstrand type (BOCD), enchondromatosis multiple (ENCHOM), Eiken skeletal dysplasia (EISD) and primary failure of tooth eruption (PFE).</p> <p>Name: PTH1R,PFE,PTHR,PTHR1</p>
Gene ID:	5745
UniProt:	<a href="#">Q03431</a>
Pathways:	<a href="#">Regulation of Carbohydrate Metabolic Process</a>

## Application Details

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.