antibodies .- online.com







RSP01 Protein (His tag)



Overview

| Quantity: | 10 μg |
|-------------------------------|--|
| Target: | RSP01 |
| Origin: | Mouse |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This RSP01 protein is labelled with His tag. |

Product Details

| Purpose: | Recombinant Mouse R-spondin-1/RSP01 Protein |
|------------------|---|
| Sequence: | SRGIKGKRQR RISAEGSQAC AKGCELCSEV NGCLKCSPKL FILLERNDIR QVGVCLPSCP |
| | PGYFDARNPD MNKCIKCKIE HCEACFSHNF CTKCQEALYL HKGRCYPACP EGSTAANSTM |
| | ECGSPAQCEM SEWSPWGPCS KKRKLCGFRK GSEERTRRVL HAPGGDHTTC SDTKETRKCT |
| | VRRTPCPEGQ KRRKGGQGRR ENANRHPARK NSKEPRSNSR RHKGQQQPQP GTTGPLTSVG |
| | PTWAQ |
| Specificity: | Ser21-Gln265 |
| Purity: | > 97 % by SDS-PAGE. |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | < 0.01EU/µg |

Target Details

| Target: | RSP01 |
|---------|-------|
| | |

Target Details

| Alternative Name: | R-spondin-1/RSP01 (RSP01 Products) |
|---------------------|--|
| Background: | Description: RSP01 gene is a member of the R-spondin family. It encodes RSP01 which is |
| | known as a secreted activator protein with two cystein-rich, furin-like domains and one |
| | thrombospondin type 1 domain. In mice, RSPO1 induces the rapid onset of crypt cell |
| | proliferation and increases intestinal epithelial healing, providing a protective effect against |
| | chemotherapy-induced adverse effects. This protein is an activator of the beta-catenin |
| | signaling cascade, leading to TCF-dependent gene activation. RSPO1 acts both in the canonica |
| | Wnt/beta-catenin-dependent pathway and in non-canonical Wnt signaling pathway, probably by |
| | acting as an inhibitor of ZNRF3, an important regulator of the Wnt signaling pathway. It also |
| | acts as a ligand for frizzled FZD8 and LRP6. |
| | Name: Rspondin, R-spondin, RSP01 |
| Gene ID: | 192199 |
| UniProt: | Q9Z132 |
| Application Details | |

Application Details

|--|

Handling

| Format: | Lyophilized |
|------------------|---|
| Reconstitution: | Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles. |
| Concentration: | 0.47 mg/mL |
| Buffer: | Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. |