

Datasheet for ABIN7520442 Chemerin Protein (Fc Tag)



Overview Quantity: 20 µg Chemerin (RARRES2) Target: Origin: Human Source: HEK-293 Cells Protein Type: Recombinant Purification tag / Conjugate: This Chemerin protein is labelled with Fc Tag. **Product Details** Recombinant Human RARRES2/TIG2 Protein Purpose: Sequence: ELTEAQRRGL QVALEEFHKH PPVQWAFQET SVESAVDTPF PAGIFVRLEF KLQQTSCRKR DWKKPECKVR PNGRKRKCLA CIKLGSEDKV LGRLVHCPIE TQVLREAEEH QETQCLRVQR AGEDPHSFYF PGQFAFS Specificity: Glu21-Ser157 Purity: > 95 % by SDS-PAGE. Sterility: 0.22 µm filtered Endotoxin Level: < 0.01EU/µg Target Details

Target:	Chemerin (RARRES2)
Alternative Name:	RARRES2/TIG2 (RARRES2 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7520442 | 07/24/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Background:	Description: Retinoic acid receptor responder protein 2 (RARRES2) is a small secreted protein
	involved in multiple cancers, including adrenocortical carcinoma (ACC). Serum RARRES2 may
	be used as a novel prognostic marker for ACC. Retinoic acid receptor responder 2 (RARRES2) is
	transcriptionally downregulated in multiple cancer types. Previous studies suggested that it can
	serve as an immune-dependent tumor suppressor by acting as a chemoattractant to recruit
	anticancer immune cells expressing its receptor, the chemerin chemokine receptor 1
	(CMKLR1), to sites of tumor. Mechanistically, RARRES2 overexpression in ACC cells inhibited
	Wnt/beta-catenin pathway activity by promoting beta-catenin phosphorylation and degradation,
	it also inhibited the phosphorylation of p38 mitogen-activated protein kinase. Thus RARRES2 is
	a novel tumor suppressor for ACC, which can function through an immune-independent
	mechanism.
	Name: TIG2, HP10433,RARRES2
Gene ID:	5919
UniProt:	Q99969
Pathways:	Brown Fat Cell Differentiation

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Concentration:	0.8 mg/mL
Buffer:	Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN7520442 | 07/24/2024 | Copyright antibodies-online. All rights reserved.