antibodies -online.com





TYRO3 Protein (Fc Tag, His tag)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	10 μg
Target:	TYRO3
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This TYRO3 protein is labelled with Fc Tag,His tag.

Product Details

Purpose:	Active Recombinant Human TYR03 Protein
Sequence:	AGLKLMGAPV KLTVSQGQPV KLNCSVEGME EPDIQWVKDG AVVQNLDQLY IPVSEQHWIG
	FLSLKSVERS DAGRYWCQVE DGGETEISQP VWLTVEGVPF FTVEPKDLAV PPNAPFQLSC
	EAVGPPEPVT IVWWRGTTKI GGPAPSPSVL NVTGVTQSTM FSCEAHNLKG LASSRTATVH
	LQALPAAPFN ITVTKLSSSN ASVAWMPGAD GRALLQSCTV QVTQAPGGWE VLAVVVPVPP
	FTCLLRDLVP ATNYSLRVRC ANALGPSPYA DWVPFQTKGL APASAPQNLH AIRTDSGLIL
	EWEEVIPEAP LEGPLGPYKL SWVQDNGTQD ELTVEGTRAN LTGWDPQKDL IVRVCVSNAV
	GCGPWSQPLV VSSHDRAGQQ GPPHSRTS
Specificity:	Ala41-Ser428
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 μm filtered
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.

Product Details

Biological Activity Comment:

Measured by its binding ability in a functional ELISA. Immobilized Human Gas6 at 2 μ g/mL (100 μ L/well) can bind Human TYRO3 with a linear range of 0.1-5.6 ng/mL.

Target Details

Target:	TYRO3
Alternative Name:	TYRO3 (TYRO3 Products)
Background:	Description: TYRO3, also known as Axl (Ufo, Ark), Dtk (Sky, Tyro3, Rse, Brt) and Mer (human and mouse homologues of chicken c-Eyk) constitute a new receptor tyrosine kinase subfamily. TYRO3 protein is activated by the products of the growth arrest-specific gene 6 and protein S genes and is involved in controlling cell survival and proliferation, spermatogenesis, immunoregulation and phagocytosis. TYRO3 has also been identified as a cell entry factor for Ebola and Marburg viruses. Name: BYK, Dtk, Etk-2, RSE, Rek, Sky, Tif,TYRO3,Dtk,Etk-2,RSE,Rek,Sky,Tif
Gene ID:	7301
UniProt:	Q06418
Pathways:	RTK Signaling

Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile
	distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is
	recommended to add a carrier protein or stablizer (e.g. 0.1 $\%$ BSA, 5 $\%$ HSA, 10 $\%$ FBS or 5 $\%$
	Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80 °C for long term.
	After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1
	week.