antibodies .- online.com





anti-Glycogenin 2 antibody (AA 401-501) (Biotin)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	Glycogenin 2 (GYG2)
Binding Specificity:	AA 401-501
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Glycogenin 2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GYG2	
Isotype:	IgG	
Predicted Reactivity:	Human,Dog	
Purification:	Purified by Protein A.	

Target Details

Target:	Glycogenin 2 (GYG2)
Alternative Name:	GYG2 (GYG2 Products)
Background:	Synonyms: Glycogenin 2, Glycogenin 2, GN 2, GN 2, GYG 2, GYG 2, OTTHUMP00000022855,

GLYG2	HIIN	$IA \Delta N$
CILICIZ	LIUI	/I/AIN.

Background: This gene encodes a member of the the glycogenin family. Glycogenin is a self-glucosylating protein involved in the initiation reactions of glycogen biosynthesis. A gene on chromosome 3 encodes the muscle glycogenin and this X-linked gene encodes the glycogenin mainly present in liver, both are involved in blood glucose homeostasis. This gene has a short version on chromosome Y, which is 3' truncated and can not make a functional protein. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

Gene ID: 8908

Pathways: Cellular Glucan Metabolic Process

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months