

Datasheet for ABIN753152

anti-Methionine Adenosyltransferase antibody (AA 31-130) (Cy5)



Go to Product page

		۱۷(e	rV	16	9	M	V
--	--	-----	---	----	----	---	---	---

Overview	
Quantity:	100 μL
Target:	Methionine Adenosyltransferase (MAT)
Binding Specificity:	AA 31-130
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Methionine Adenosyltransferase antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence
	(Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human Methionine adenosyltransferase
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Dog, Cow, Pig, Chicken, Rabbit, Zebrafish, Monkey
Purification:	Purified by Protein A.
Target Details	
Target Details Target:	Methionine Adenosyltransferase (MAT)

Target Details

9				
Background:	Synonyms: MAT2A AdoMet synthetase 2, AdoMet synthetase, AMS 2, AMS2, MAT 2A, MAT II,			
	MATA 2, MATA2, MATII, Methionine adenosyltransferase 2, Methionine adenosyltransferase,			
	Methionine adenosyltransferase II alpha, Methionine adenosyltransferase II, S			
	adenosyl methion in esynthetasegammaform, Sadenosyl methion in esynthetaseiso formtype			
	2, SAMS 2, SAMS2, METK1_HUMAN.			
	Background: S adenosylmethionine synthetase catalyzes the formation of S			
	adenosylmethionine from methionine and ATP. In mammalian tissues, there are three distinct			
	forms of AdoMet synthases designated as alpha, beta, and gamma. Alpha and beta are			
	expressed only in adult liver, while gamma is widely distributed in extrahepatic tissues.			
Gene ID:	4143			
Application Details				
Application Notes:	IF(IHC-P) 1:50-200			
	IF(IHC-F) 1:50-200			
	IF(ICC) 1:50-200			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	1 μg/μL			
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and			
	50 % Glycerol.			
Preservative:	ProClin			
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be			
	handled by trained staff only.			
Storage:	-20 °C			
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.			
Expiry Date:	12 months			