

Datasheet for ABIN753187 anti-SMURF2 antibody (AA 601-700) (HRP)



\sim			
()\	/ e	rVI	iew

Overview	
Quantity:	100 μL
Target:	SMURF2
Binding Specificity:	AA 601-700
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SMURF2 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human SMURF2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	SMURF2
Alternative Name:	SMURF2 (SMURF2 Products)

Target Details

Background:	Synonyms: E3 ubiquitin-protein ligase SMURF2, hSMURF2, SMAD ubiquitination regulatory
	factor 2, SMAD-specific E3 ubiquitin-protein ligase 2, SMURF2
	Background: E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-
	conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to
	targeted substrates. Interacts with SMAD1 and SMAD7 in order to trigger their ubiquitination
	and proteasome-dependent degradation. In addition, interaction with SMAD7 activates
	autocatalytic degradation, which is prevented by interaction with SCYE1. Forms a stable
	complex with the TGF-beta receptor-mediated phosphorylated SMAD2 and SMAD3. In this way,
	SMAD2 may recruit substrates, such as SNON, for ubiquitin-mediated degradation. Enhances
	the inhibitory activity of SMAD7 and reduces the transcriptional activity of SMAD2.
	Coexpression of SMURF2 with SMAD1 results in considerable decrease in steady-state level of
	SMAD1 protein and a smaller decrease of SMAD2 level.
Gene ID:	64750
UniProt:	Q9HAU4
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish
	peroxidase.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months