# antibodies .- online.com





# Datasheet for ABIN7533705

# **MANF Protein**



#### Overview

Quantity:	100 μg
Target:	MANF
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant

#### **Product Details**

Purpose:	Recombinant Human MANF Protein
Sequence:	LRPGDCEVCI SYLGRFYQDL KDRDVTFSPA TIENELIKFC REARGKENRL CYYIGATDDA ATKIINEVSK PLAHHIPVEK ICEKLKKKDS QICELKYDKQ IDLSTVDLKK LRVKELKKIL DDWGETCKGC AEKSDYIRKI NELMPKYAPK AASARTDL
Specificity:	Leu25-Leu182
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.

## **Target Details**

Target:	MANF
Alternative Name:	MANF (MANF Products)
Background:	Description: The protein is localized in the endoplasmic reticulum (ER) and golgi, and is also
	secreted. Reducing expression of this protein increases susceptibility to ER stress-induced

#### **Target Details**

death and results in cell proliferation. Activity of this protein is important in promoting the survival of dopaminergic neurons. The presence of polymorphisms in the N-terminal arginine-rich region, including a specific mutation that changes an ATG start codon to AGG, have been reported in a variety of solid tumors, however, these polymorphisms were later shown to exist in Normal tissues and are thus no longer thought to be tumor-related.

Gene ID:

7873

UniProt:

P55145

## **Application Details**

Restrictions:

For Research Use only

Name: MANF, ARMET, ARP

## Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80 °C for long term.  After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.