



[Go to Product page](#)

## Datasheet for ABIN7533732 PPIA Protein

### Overview

Quantity:	100 µg
Target:	PPIA
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant

### Product Details

Purpose:	Recombinant Human PPIase A/PPIA Protein
Sequence:	MVNPTVFFDI AVDGEPLGRV SFELFADKVP KTAENFRALS TGEKGFYK SCFHRIIPGF MCQGGDFTRH NGTGGKSIYG EK FEDENFIL KHTGPGILSM ANAGPNTNGS QFFICTAKTE WLDGKHVVFG KVKEGMNIVE AMERFGSRNG KTSKKITIAD CGQLE
Specificity:	Met1-Glu165
Purity:	> 97 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	< 1.0 EU/µg of the protein by LAL method.

### Target Details

Target:	PPIA
Alternative Name:	PPIase A/PPIA ( <a href="#">PPIA Products</a> )
Background:	Description: This protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides

## Target Details

---

and accelerate the folding of proteins. The protein is a cyclosporin binding-protein and may play a role in cyclosporin A-mediated immunosuppression. The protein can also interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions.

Name: PPIA,CYPA,CYPH,HEL-S-69p

---

Gene ID: 5478

---

UniProt: [P62937](#)

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

---

Reconstitution: Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

---

Buffer: Lyophilized from a 0.22 µm filtered solution of 50 mM Tris, 150 mM NaCl, pH 8.0.

---

Storage: -20 °C,-80 °C

---

Storage Comment: Store the lyophilized protein at -20°C to -80 °C for long term.  
After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.