



[Go to Product page](#)

Datasheet for ABIN7533766
Growth Hormone 1 Protein (GH1) (His tag)

Overview

Quantity:	100 µg
Target:	Growth Hormone 1 (GH1)
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This Growth Hormone 1 protein is labelled with His tag.

Product Details

Purpose:	Active Recombinant Human Somatotropin/GH-N/GH1 Protein
Sequence:	FPTIPLSRLF DNAMLRAHRL HQLAFDITYQE FEEAYIPKEQ KYSFLQNPQT SLCFSESIPT PSNREETQK SNLELLRISL LLIQSWLEPV QFLRSVFANS LVIYASDSNV YDLLKDLEEG IQTLMGRLED GSPRTGQIFK QTYSKFDTNS HNDDALLKNY GLLYCFRKDM DKVETFLRIV QCRSVEGSCG F
Specificity:	Phe27-Phe217
Purity:	> 92 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized Human GH1 at 2 µg/mL (100 µL/well) can bind Human GHR with a linear range of 0.1-19.4 ng/mL.

Target Details

Target: Growth Hormone 1 (GH1)

Alternative Name: Somatotropin/GH-N/GH1 ([GH1 Products](#))

Background: Description: Growth hormone (GH), also known as somatotropin, is a member of a family of growth factors that includes prolactin, placental lactogens, proliferins, and somatolactin. It is synthesized primarily by somatotropes in the anterior pituitary and is stored in secretory granules. The pulsatile release of GH into circulation is regulated by the concerted actions of the hypothalamic hormones - GH-releasing hormone (GHRH) and somatostatin (SST) - as well as by signals from the periphery - ghrelin and leptin. Human GH is a pleiotropic cytokine that exerts its biological actions by binding to the transmembrane GH receptor, which is present in many cell types. GH stimulates the liver and other tissues to produce IGF-1, which regulates growth and metabolism. GH has also been shown to have direct effects on growth that is independent of IGF-1. GH, directly or indirectly via IGF-1, can act on B cells, T cells, NK cells, macrophages and neutrophils to exert immunomodulatory activities. In addition, GH can act directly on various cell types to induce lipolysis, lactation, amino acid uptake and protein synthesis.

Name: GH1, GH, GH-N, GHB5, GHN, IGHD1B, hGH-N, somatotropin,GH,GH-N,GHB5,GHN,IGHD1B,hGH-N

Gene ID: 2688

UniProt: [P01241](#)

Pathways: [NF-kappaB Signaling](#), [JAK-STAT Signaling](#), [Intracellular Steroid Hormone Receptor Signaling Pathway](#), [Peptide Hormone Metabolism](#), [Regulation of Intracellular Steroid Hormone Receptor Signaling](#), [Regulation of Hormone Metabolic Process](#), [Response to Growth Hormone Stimulus](#), [Regulation of Hormone Biosynthetic Process](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Handling

Buffer: Lyophilized from a 0.22 µm filtered solution of 20 mM Tris, 150 mM NaCl, pH 8.0.

Storage: -20 °C,-80 °C

Storage Comment: Store the lyophilized protein at -20°C to -80 °C for long term.
After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.