

Datasheet for ABIN7533801  
**CSTA Protein (His tag)**



[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	CSTA
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CSTA protein is labelled with His tag.

## Product Details

Purpose:	Active Recombinant Human Cystatin-A/CSTA Protein
Sequence:	MIPGGLSEAK PATPEIQEIV DKVKPQLEEK TNETYGKLEA VQYKTQVVAG TNYIYKVRAG DNKYMHLKVF KSLPGQNEDL VLTGYQVDKN KDELDTGF
Specificity:	Met1-Phe98
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	< 1.0 EU/µg of the protein by LAL method.
Biological Activity Comment:	Measured by its ability to inhibit papain cleavage of a fluorogenic peptide substrate Z-FR-AMC. The IC50 value is <0.7 nM.

## Target Details

Target:	CSTA
Alternative Name:	Cystatin-A/CSTA ( <a href="#">CSTA Products</a> )
Background:	<p>Description: The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins, and kininogens. This gene encodes a stefin that functions as a cysteine protease inhibitor, forming tight complexes with papain and the cathepsins B, H, and L. The protein is one of the precursor proteins of cornified cell envelope in keratinocytes and plays a role in epidermal development and maintenance. Stefins have been proposed as prognostic and diagnostic tools for cancer.</p> <p>Name: CSTA,AREI,PSS4,STF1,STFA</p>
Gene ID:	1475
UniProt:	<a href="#">P01040</a>
Pathways:	<a href="#">Response to Water Deprivation</a>

## Application Details

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 µm filtered solution of 20 mM Tris, 150 mM NaCl, pH 8.0.
Storage:	-20 °C,-80 °C
Storage Comment:	<p>Store the lyophilized protein at -20°C to -80 °C for long term.</p> <p>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.</p>