



[Go to Product page](#)

Datasheet for ABIN7534161
IL12A Protein (Fc Tag,His tag)

Overview

Quantity:	100 µg
Target:	IL12A
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL12A protein is labelled with Fc Tag,His tag.

Product Details

Purpose:	Recombinant Human IL-12A Protein
Sequence:	RNLPVATPDP GMFPC LHHSQ NLLRAVSNML QKARQTLEFY PCTSEEIDHE DITKDKTSTV EACLPLELTK NESCLNSRET SFITNGSCLA SRKTSFMMAL CLSSIYEDLK MYQVEFKTMN AKLLMDPKRQ IFLDQNMLAV IDELMQALNF NSETVPQKSS LEEPDFYKTK IKLCILLHAF RIRAVTIDRV MSYLNAS
Specificity:	Arg23-Ser219
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.

Target Details

Target:	IL12A
Alternative Name:	IL-12A (IL12A Products)

Target Details

Background: Description: Interleukin-12 subunit alpha (IL12A/IL-12p35) is also known as Cytotoxic lymphocyte maturation factor 35 kDa subunit, cytotoxic lymphocyte maturation factor 1, p35, NK cell stimulatory factor chain 1, and interleukin-12 alpha chain. IL12A/IL-12p35 is a subunit of a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. The cytokine is a disulfide-linked heterodimer composed of the 35-kD subunit encoded by this gene, and a 40-kD subunit that is a member of the cytokine receptor family. This cytokine is required for the T-cell-independent induction of interferon (IFN)-gamma, and is important for the differentiation of both Th1 and Th2 cells. The responses of lymphocytes to this cytokine are mediated by the activator of transcription protein STAT4. Nitric oxide synthase 2A (NOS2A/NOS2) is found to be required for the signaling process of this cytokine in innate immunity.

Name: P35, CLMF, NFSK, NKSF1, IL-12A,IL12A

Gene ID: 3592

UniProt: [P29459](#)

Pathways: [JAK-STAT Signaling](#), [Regulation of Leukocyte Mediated Immunity](#), [Positive Regulation of Immune Effector Process](#), [Activated T Cell Proliferation](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Buffer: Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Storage: -20 °C,-80 °C

Storage Comment: Store the lyophilized protein at -20°C to -80 °C for long term.
After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.