# antibodies .- online.com





## Datasheet for ABIN7534276

# **LUM Protein (His tag)**



#### Overview

Quantity:	100 μg
Target:	LUM
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This LUM protein is labelled with His tag.

#### **Product Details**

Purpose:	Recombinant Human Lumican/LUM Protein
Sequence:	QYYDYDFPLS IYGQSSPNCA PECNCPESYP SAMYCDELKL KSVPMVPPGI KYLYLRNNQI
	DHIDEKAFEN VTDLQWLILD HNLLENSKIK GRVFSKLKQL KKLHINHNNL TESVGPLPKS
	LEDLQLTHNK ITKLGSFEGL VNLTFIHLQH NRLKEDAVSA AFKGLKSLEY LDLSFNQIAR
	LPSGLPVSLL TLYLDNNKIS NIPDEYFKRF NALQYLRLSH NELADSGIPG NSFNVSSLVE
	LDLSYNKLKN IPTVNENLEN YYLEVNQLEK FDIKSFCKIL GPLSYSKIKH LRLDGNRISE
	TSLPPDMYEC LRVANEVTLN
Specificity:	Gln19-Asn338
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 μm filtered
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.

## **Target Details**

Target:	LUM
Alternative Name:	Lumican/LUM (LUM Products)
Background:	Description: Lumican is a member of the family of small leucine-rich repeat proteoglycans
	(SLRPs) and the class II subfamily .which is a major component of the cornea, dermal, and
	muscle connective tissues. Lumican is the major keratan sulfate proteoglycan of the cornea but
	is also distributed in interstitial collagenous matrices throughout the body. Lumican regulates
	collagenous matrix assembly as a keratan sulfate proteoglycan in the cornea and is also
	present in the connective tissues of other organs and embryonic corneal stroma as a
	glycoprotein. Lumican may regulate collagen fibril organization and circumferential growth,
	corneal transparency, and epithelial cell migration and tissue repair. Lumican's over-expression
	has been reported in carcinoid tumor, breast, colorectal, neuroendocrine, uterine cervical and
	pancreatic cancers
	Name: LDC, SLRR2D,LUM,SLRR2D,lumican
Gene ID:	4060
UniProt:	P51884
Pathways:	Glycosaminoglycan Metabolic Process
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile
	distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is
	recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 %
	Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80 °C for long term.
	After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1
	week.