



[Go to Product page](#)

Datasheet for ABIN7534330
ULBP3 Protein (Fc Tag,His tag)

Overview

Quantity:	100 µg
Target:	ULBP3
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ULBP3 protein is labelled with Fc Tag,His tag.

Product Details

Purpose:	Active Recombinant Human NKG2D ligand 3/ULBP3 Protein
Sequence:	GRADAHSLWY NFTIIHLPRH GQQWCEVQSQ VDQKNFLSYD CGSDKVLSMG HLEEQLYATD AWGKQLEMLR EVGQRLRLEL ADTELEDFTP SGPLTLQVRM SCECEADGYI RGSWQFSFDG RKFLFDSNN RKWTVVHAGA RRMKEKWEKD SGLTTFKVM SMRDCKSWLR DFLMHRKKRL EPTAPPTMAP
Specificity:	Gly27-Pro216
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.01 EU/µg of the protein by LAL method.

Target Details

Target:	ULBP3
Alternative Name:	NKG2D ligand 3/ULBP3 (ULBP3 Products)

Target Details

Background: Description: ULBP-3 is a member of a family of cell-surface proteins that function as ligands for human NKG2D. ULBP-3 has also been described under the names RaeT1N (retinoic acid early transcript), NKG2DL3, and ALCAN-gamma. The name ULBP-3 derives from the original identification of three proteins, ULBP-1, -2, and -3, as ligands for the human cytomegalovirus glycoprotein UL16, they were designated UL16 binding proteins (ULBP). The gene for ULBP-3 resides in a cluster of ten related genes, six of which encode potentially functional glycoproteins. Amino acid sequence identity within this family ranges from 30-60 % . These proteins are distantly related to MHC class I proteins, but they possess only the alpha 1 and alpha 2 Ig-like domains, and they have no capacity to bind peptide or interact with beta 2-microglobulin. Some family members, including ULBP-3, are anchored to the membrane via a GPI-linkage, whereas others have transmembrane domains.

Name: ULBP3

Gene ID: 79465

UniProt: [Q9BZM4](#)

Pathways: [Regulation of Leukocyte Mediated Immunity](#), [Positive Regulation of Immune Effector Process](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Buffer: Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Storage: -20 °C,-80 °C

Storage Comment: Store the lyophilized protein at -20°C to -80 °C for long term.
After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.