# antibodies .- online.com





# **CDH6 Protein (His tag)**



## Overview

Quantity:	100 μg
Target:	CDH6
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CDH6 protein is labelled with His tag.

# **Product Details**

Purpose:	Active Recombinant Human CDH6/K-Cadherin Protein
Sequence:	TLSTPLSKRT SGFPAKKRAL ELSGNSKNEL NRSKRSWMWN QFFLLEEYTG SDYQYVGKLH
	SDQDRGDGSL KYILSGDGAG DLFIINENTG DIQATKRLDR EEKPVYILRA QAINRRTGRP
	VEPESEFIIK IHDINDNEPI FTKEVYTATV PEMSDVGTFV VQVTATDADD PTYGNSAKVV
	YSILQGQPYF SVESETGIIK TALLNMDREN REQYQVVIQA KDMGGQMGGL SGTTTVNITL
	TDVNDNPPRF PQSTYQFKTP ESSPPGTPIG RIKASDADVG ENAEIEYSIT DGEGLDMFDV
	ITDQETQEGI ITVKKLLDFE KKKVYTLKVE ASNPYVEPRF LYLGPFKDSA TVRIVVEDVD
	EPPVFSKLAY ILQIREDAQI NTTIGSVTAQ DPDAARNPVK YSVDRHTDMD RIFNIDSGNG
	SIFTSKLLDR ETLLWHNITV IATEINNPKQ SSRVPLYIKV LDVNDNAPEF AEFYETFVCE
	KAKADQLIQT LHAVDKDDPY SGHQFSFSLA PEAASGSNFT IQDNKDNTAG ILTRKNGYNR
	HEMSTYLLPV VISDNDYPVQ SSTGTVTVRV CACDHHGNMQ SCHAEALIHP TGLSTGA
Specificity:	Thr19-Ala615
Purity:	> 95 % by SDS-PAGE.

#### **Product Details**

Product Details	
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized Human CDH6 at $1\mu g/mL$ (100 $\mu L/well$ ) can bind CDH6 Mouse mAb with a linear range of 0.05-6.31 ng/mL.
Target Details	
Target:	CDH6
Alternative Name:	CDH6/K-Cadherin (CDH6 Products)

#### Background:

Description: CDH6 is a family of calcium-dependent, cell-cell adhesion molecules that play an important morphoregulatory role in a wide variety of tissues. Alterations in cadherin function have been implicated in tumor progression in a number of adenocarcinomas. Cadherin-6 (CDH6), also known as K-cadherin (KCAD), is a type-II classic cadherin cell-cell adhesion molecules, which are expressed in graded or areal patterns, as well as layer-specific patterns, in the cortical plate. Human Cadherin-6 is synthesized as a 790 aa type I transmembrane glycoprotein that contains a 18 aa signal peptide, a 35 aa propeptide, a 562 aa extracellular region, a 21 aa transmembrane segment, and a 154 aa cytoplasmic domain. There are five cadherin domains of approximately 110 aa each in the extracellular region. Cadherin-6 is highly expressed in brain, cerebellum, and kidney, and may contribute to the formation of the segmental structure of the early brain, as well as the development of renal proximal tubules. Weak expression is also detected lung, pancreas, and gastric mucosa. Additionally, it is specifically expressed in the proximal tubule of normal kidneys and in renal cell cancer. Thus, Cadherin-6 is a new prognostic factor for renal cancer.

Name: CAD6, KCAD, CDH6, KCAD

Gene ID:	1004
UniProt:	P55285-1
Pathways:	Cell-Cell Junction Organization

### **Application Details**

Restrictions:	For Research Use only	

## Handling

Format: Lyophilized
---------------------

# Handling

Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile	
	distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is	
	recommended to add a carrier protein or stablizer (e.g. 0.1 $\%$ BSA, 5 $\%$ HSA, 10 $\%$ FBS or 5 $\%$	
	Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.	
Buffer:	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Store the lyophilized protein at -20°C to -80 °C for long term.	
	After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1	
	week.	