

Datasheet for ABIN7535358
VSIG4 Protein (His tag)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	VSIG4
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This VSIG4 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human VSIG4 Protein
Sequence:	RPILVPEVS TGPWKGDVNL PCTYDPLQGY TQVLVKWLVQ RGSDPVTIFL RDSSGDHIQQ AKYQGRLHVS HKVPGDVSLQ LSTLEMDDRS HYTCEVTWQT PDGNQVVRDK ITELRVQKLS VSKPTVTTGS GYGFTVPQGM RISLQCQARG SPPISYIWYK QQTNNQEPIK VATLSTLLFK PAVIADSGSY FCTAKGQVGS EQHSDIVKFV VKDSSKLLKT KTEAPTTMTY PLKATSTVKQ SWDWTTDMDG YLGETSAGPG KSLP
Specificity:	Arg20-Pro283
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	<0.1EU/µg

Target Details

Target:	VSIG4
---------	-------

Target Details

Alternative Name: VSIG4 ([VSIG4 Products](#))

Background: Description: VSIG4 (V-set and immunoglobulin domain containing 4), also known as complement receptor of the immunoglobulin superfamily (CRIg) and Z39Ig, is a type I transmembrane glycoprotein. It is a B7 family-related protein and an Ig superfamily member. In contrast to the B7 family members which contain two IgG domains, VSIG4 contains one complete V-type Ig domain and a truncated C-type Ig domain. VSIG4 is exclusively expressed on tissue resident macrophages and binds to multimers of C3b and iC3b that are covalently attached to particle surfaces. No VSIG4 expression appears to be present in T and B cells. VSIG4 functions as a negative regulator of T cell activation, and may be involved in the maintenance of peripheral T cell tolerance, and is also identified as a potent suppressor of established inflammation. Mouse VSIG4 is synthesized as a 28 amino acid precursor that contains a signal sequence, a V-type Ig domain (aa 36-115), one potential N-linked glycosylation site, and a single transmembrane domain. The V-type Ig domain of mouse VSIG4 shares 86 % and 8 % aa sequence identity with the V-type Ig domains of rat and human VSIG4, respectively.

Name: CRIg,Z39IG,VSIG4,VSIG4,CRIg,Z39IG,VSIG4

Gene ID: 11326

UniProt: [Q9Y279-1](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Buffer: Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Storage: -20 °C,-80 °C

Storage Comment: Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.