



[Go to Product page](#)

Datasheet for ABIN7535659
CPE Protein (Fc Tag,His tag)

Overview

Quantity:	100 µg
Target:	CPE
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CPE protein is labelled with Fc Tag,His tag.

Product Details

Purpose:	Recombinant Human Carboxypeptidase E/CPE Protein
Sequence:	MAGRGGSSALL ALCGALAACG WLLGAEAQEP GAPAAAGMRRR RRLQQEDGIS FEYHRYPELR EALVSVWLQC TAISRIYTVG RSFEGRELLV IELSDNPGVH EPGEPEFKYI GNMHGNEAVG RELLIFLAQY LCNEYQKGNE TIVNLIHSTR IHIMP SLNPD GFKAASQPG ELKDWVFGRS NAQGIDLNRN FPDLDRIYVY NEKEGGPNNH LLKNMKKIVD QNTKLAPETK AVIHWIMDIP FVLSANLHGG DLVANYPYDE TRSGSAHEYS SSPDDAIFQS LARAYSSFNP AMSDPNRPPC RKNDDDSSFV DGT TNGGAWY SVPGGMQDFN YLSSNCFEIT VELSCEKFPP EETLKTYWED NKNSLISYLE QIHRGVKGFV RDLQGNPIAN ATISVEGIDH DV TSAKDGDY WRLLIPGNYK LTASAPGYLA ITKKVAVPYS PAAGVDFELE SFS
Specificity:	Met1-Ser453
Purity:	> 85 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	<0.1EU/µg

Target Details

Target:	CPE
Alternative Name:	Carboxypeptidase E/CPE (CPE Products)
Background:	<p>Description: Carboxypeptidase E (CPE), also known as Carboxypeptidase H, is a peripheral membrane protein and a zinc metalloprotein, and the conversion of proCPE into CPE occurs primarily in secretory vesicles. The active form of CPE cleaves C-terminal amino acid residues of the peptide, and is thus involved in the biosynthesis of peptide hormones and neurotransmitters including insulin, enkephalin, etc. The enzymatic activity is enhanced by millimolar concentrations of Co²⁺. It has also been proposed that membrane-associated carboxypeptidase E acts as a sorting receptor for targeting regulated secretory proteins which are mostly prohormones and neuropeptides in the trans-Golgi network of the pituitary and in secretory granules into the secretory pathway. Its interaction with glycosphingolipid-cholesterol rafts at the TGN facilitates the targeting. Mutations in this gene are implicated in type II diabetes due to impaired glucose clearance and insulin resistance.</p> <p>Name: CPE,CPH</p>
Gene ID:	1363
UniProt:	P16870-1
Pathways:	Peptide Hormone Metabolism , Synaptic Membrane

Application Details

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
Storage:	-20 °C, -80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.