

## Datasheet for ABIN7535778

# **IL18BP Protein (His tag)**



## Overview

Alternative Name:

Quantity:	100 μg
Target:	IL18BP
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL18BP protein is labelled with His tag.
Product Details	
Purpose:	Recombinant Mouse IL-18BP Protein
Sequence:	TSAPQTTATV LTGSSKDPCS SWSPAVPTKQ YPALDVIWPE KEVPLNGTLT LSCTACSRFP YFSILYWLGN GSFIEHLPGR LKEGHTSREH RNTSTWLHRA LVLEELSPTL RSTNFSCLFV DPGQVAQYHI ILAQLWDGLK TAPSPSQETL SSHSPVSRSA GPGVA
Specificity:	Thr29-Ala193
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	<0.1EU/µg
Target Details	
Target:	IL18BP

IL-18BP (IL18BP Products)

#### **Target Details**

Bac	kar	'nΙ	ın	Ч.
Duo		$\sim$	<i>.</i>	ч.

Description: Interleukin-18-binding protein (IL-18BP) is a constitutively expressed and secreted protein. IL-18BP is a cytokine receptor that belongs to the interleukin 1 receptor family. This receptor specifically binds interleukin 18 (IL18) and is essential for IL18 mediated signal transduction. IFN-alpha and IL12 are reported to induce the expression of this receptor in NK and T cells. This gene along with four other members of the interleukin 1 receptor family, including IL1R2, IL1R1, ILRL2 (IL-1Rrp2), and IL1RL1 (T1/ST2), form a gene cluster on chromosome 2q. The adjacently located family members IL18 Receptor 1 (IL18R1) and IL18 receptor accessory protein (IL18RAP) may also be important in the development of asthma and atopy. IL-18 binding protein (IL-18BP) was only moderately elevated, resulting in a high level of biologically active free IL-18 in HPS. A severe IL-18/IL-18BP imbalance results in Th-1 lymphocyte and macrophage activation, which escapes control by NK-cell cytotoxicity and may allow for secondary HPS in patients with underlying diseases.

Name: MC54L, Igifbp, IL-18BP, IL18BP

Gene ID:

16068

UniProt:

Q9Z0M9

## **Application Details**

Restrictions:

For Research Use only

## Handling

Format:	Lyophilized		
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile		
	distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is		
	recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 %		
	Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.		
Buffer:	Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.		
Storage:	-20 °C,-80 °C		
Storage Comment:	Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein		
	solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.		