

# Datasheet for ABIN7535965

# **CXCL7 Protein**



#### Overview

| Quantity:    | 100 μg                     |
|--------------|----------------------------|
| Target:      | CXCL7 (PPBP)               |
| Origin:      | Human                      |
| Source:      | Escherichia coli (E. coli) |
| Protein Tyne | Recombinant                |

#### **Product Details**

| Purpose:     | Recombinant Human CXCL7/PPBP Protein  |
|--------------|---|
| Sequence:    | AELRCMCIKT TSGIHPKNIQ SLEVIGKGTH CNQVEVIATL KDGRKICLDP DAPRIKKIVQ<br>KKLAGDESAD |
| Specificity: | Ala59-Asp128  |
| Purity:      | > 95 % by SDS-PAGE.   |
| Sterility:   | 0.22 µm filtered  |

# **Target Details**

| Target:           | CXCL7 (PPBP)   |
|-------------------|--|
| Alternative Name: | CXCL7/PPBP (PPBP Products)   |
| Background:       | Description: This protein is a platelet-derived growth factor that belongs to the CXC chemokine  |
|                   | family. This growth factor is a potent chemoattractant and activator of neutrophils. It has been |
|                   | shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis,      |
|                   | intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid    |

## **Target Details**

and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by synovial cells. The protein also is an antimicrobial protein with bactericidal and antifungal activity.

Name: PPBP,B-TG1,Beta-TG,CTAP-III,CTAP3,CTAPIII,CXCL7,LA-PF4,LDGF,MDGF,NAP-

Gene ID: 5473

UniProt: P02775

## **Application Details**

Restrictions: For Research Use only

## Handling

| Format:          | Lyophilized  |
|------------------|--|
| Reconstitution:  | Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile  |
|                  | distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is   |
|                  | recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 %      |
|                  | Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.         |
| Buffer:          | Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.                                     |
| Storage:         | -20 °C,-80 °C  |
| Storage Comment: | Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein |
|                  | solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.                             |

2,PBP,SCYB7,TC1,TC2,TGB,TGB1,THBGB,THBGB1