



[Go to Product page](#)

Datasheet for ABIN7536033  
**CD63 Protein (CD63) (His tag)**

### Overview

Quantity:	100 µg
Target:	CD63
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD63 protein is labelled with His tag.

### Product Details

Purpose:	Recombinant Human CD63 Protein
Sequence:	AGYVFRDKVM SEFNNNFRQQ MENYPKNNHT ASILDRMQAD FKCCGAANYT DWEKIPSMK NRVPDSCCIN VTVGCGINFN EKAIHKEGCV EKIGGWLRKN V
Specificity:	Ala103-Val203
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	<0.1EU/µg

### Target Details

Target:	CD63
Alternative Name:	CD63 ( <a href="#">CD63 Products</a> )
Background:	Description: The cluster of differentiation (CD) system is commonly used as cell markers in

## Target Details

---

Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 63 (CD63) is a member of the CD family and the transmembrane 4 superfamily, also known as the tetraspanin family. CD63 is a cell surface glycoprotein characterized by the presence of four hydrophobic domains. CD63 had functions in mediating signal transduction processes and then regulate a variety of cellular processes such as cell proliferation, activation and motility. It has been reported that CD63 protein associated with tumor progression and served as a blood platelet activation marker and the deficiency of this protein may be associated with Hermansky-Pudlak syndrome.

Name: MLA1, ME491, LAMP-3, OMA81H, TSPAN30,CD63

---

Gene ID: 967

---

UniProt: [P08962](#)

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

---

Buffer: Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

---

Storage: -20 °C,-80 °C

---

Storage Comment: Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.