



[Go to Product page](#)

Datasheet for ABIN7536423  
**TFF1 Protein (pSer2) (His tag)**

### Overview

Quantity:	100 µg
Target:	TFF1
Protein Characteristics:	pSer2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TFF1 protein is labelled with His tag.

### Product Details

Purpose:	Recombinant Human BCEI/PS2/TFF1 Protein
Sequence:	EAQTETCTVA PRERQNCGFP GVTPSQCANK GCCFDDTVRG VPWCFYPNTI DVPPEEECEF
Specificity:	Glu25-Phe84
Purity:	> 97 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.001 EU/µg

### Target Details

Target:	TFF1
Alternative Name:	BCEI//TFF1 ( <a href="#">TFF1 Products</a> )
Background:	Description: Trefoil Factor 1 (TFF1), also known as pS2, is one of three structurally related

## Target Details

---

secreted proteins that contain trefoil domains. These domains adopt a three-leaved conformation held together by conserved intrachain disulfide bonds. TFF1 is an approximately 7 kDa peptide that plays an important role in epithelial regeneration and wound healing (1). Mature human TFF1 shares 67 % amino acid sequence identity with mouse and rat TFF1. It is expressed by goblet cells of the gastric and intestinal mucosa and by conjunctival goblet cells (2-5). TFF1 is a copper-binding protein that can form disulfide-linked homodimers, associate into disulfide-linked complexes with Gastrokine 2, and form non-covalent complexes with the mucin MUC5AC (4, 6-8). Copper enhances TFF1 homodimerization as well as its ability to promote epithelial cell motility, wound healing, and the colonization of *H. pylori* in stomach and colon epithelia (9, 10). TFF1 is down-regulated during the progression from gastritis to gastric dysplasia to gastric cancer, although it is up-regulated in breast and prostate cancers (11-13). TFF1 inhibits the formation of calcium oxalate crystals, and its excretion in the urine is reduced in patients with kidney stones (14).

Name: pS2, BCEI, HPS2, HP1.A, pNR-2, D21S21, TFF1

---

Gene ID: 7031

---

UniProt: [P04155](#)

---

Pathways: [EGFR Signaling Pathway](#)

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

---

Concentration: 1.1 mg/mL

---

Buffer: Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

---

Storage: -20 °C, -80 °C

---

Storage Comment: Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.