



[Go to Product page](#)

Datasheet for ABIN7536435
ACVA Protein (His tag)

Overview

Quantity:	100 µg
Target:	ACVA
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ACVA protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human Activin A/INHBA Protein
Sequence:	SPTPGSEGHS AAPDCPSCAL AALPKDVPNS QPEMVEAVKK HILNMLHLKK RPDVTQPVPK AALLNAIRKL HVGKVGNGY VEIEDDIGRR AEMNELMEQT SEITFAESG TARKTLHFEI SKEGSDLSVV ERAEVLWFLK VPKANRTRTK VTIRLFQQQK HPQGSLDTGE EAEEVGLKGE RSELLLSEKV VDARKSTWHV FPVSSSIQRL LDQGKSSLDV RIACEQCQES GASLVLLGKK KKKEEEGEGK KKGEGEGGAG ADEEKEQSHR PFLMLQARQS EDHPHRRRRR GLECDGKVNI CCKKQFFVSF KDIGWNDWII APSGYHANYC EGECPSHIAG TSGSSLSFHS TVINHYRMRG HSPFANLKSC CVPTKLRPMS MLYYDDGQNI IKKDIQNMIV EECGCS
Specificity:	Ser21-Ser426
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.01EU/µg

Target Details

Target:	ACVA
Alternative Name:	Activin A/INHBA (ACVA Products)
Background:	<p>Description: The inhibin beta A subunit joins the alpha subunit to form a pituitary FSH secretion inhibitor. Inhibin has been shown to regulate gonadal stromal cell proliferation negatively and to have tumor-suppressor activity. In addition, serum levels of inhibin have been shown to reflect the size of granulosa-cell tumors and can therefore be used as a marker for primary as well as recurrent disease. Because expression in gonadal and various extragonadal tissues may vary severalfold in a tissue-specific fashion, it is proposed that inhibin may be both a growth/differentiation factor and a hormone. Furthermore, the beta A subunit forms a homodimer, activin A, and also joins with a beta B subunit to form a heterodimer, activin AB, both of which stimulate FSH secretion. Finally, it has been shown that the beta A subunit mRNA is identical to the erythroid differentiation factor subunit mRNA and that only one gene for this mRNA exists in the human genome.</p> <p>Name: EDF, FRP,INHBA</p>
Gene ID:	3624
UniProt:	P08476
Pathways:	Hormone Transport , Peptide Hormone Metabolism

Application Details

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Concentration:	0.75 mg/mL
Buffer:	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein

solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.