.-online.com antibodies

Datasheet for ABIN7536684 anti-Factor VIIIC antibody



Overview

| Quantity: | 5 mg |
|--------------|--|
| Target: | Factor VIIIC |
| Reactivity: | Dog |
| Host: | Sheep |
| Clonality: | Polyclonal |
| Conjugate: | This Factor VIIIC antibody is un-conjugated |
| Application: | ELISA, Western Blotting (WB), Neutralization (Neut), Immunoprecipitation (IP), Immunoelectrophoresis (IEP), Immunodepletion (IDe) |

Product Details

| Purpose: | Canine Factor VIII Polyclonal Antibody |
|---------------|--|
| Immunogen: | Recombinant canine FVIII (cFVIII) |
| Isotype: | IgG |
| Specificity: | Canine Factor VIII |
| Purification: | Fractionation of IgG from antiserum |

Target Details

| Target: | Factor VIIIC |
|-------------------|---------------------------------------|
| Alternative Name: | Factor VIII:C (Factor VIIIC Products) |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7536684 | 05/10/2024 | Copyright antibodies-online. All rights reserved.

| Application Details | |
|---------------------|--|
| Application Notes: | Immunoblot 5-25 µg/mL, ELISA capture 20-50 µg/mL |
| Comment: | Whole IgG is a base level product and refers to IgG antibodies purified from antiserum using multiple precipitation and dialysis steps. The purity of the IgG is typically 90% and is provided in a solution of HEPES buffered saline containing 50% glycerol (v/v). The titre is essentially the same as the starting antiserum and each vial typically contains the amount of IgG recovered from one millilitre of antiserum. This level of product is generally intended for use in applications such as immunoprecipitation, immunoelectrophoresis, immuno-depletion and activity neutralization assays. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Reconstitution: | Allow product to warm to room temperature and gently mix before use |
| Concentration: | 5 mg/mL |
| Buffer: | 10 mM HEPES, pH 7.4, 150 mM NaC, 50 % (v/v) glycerol |
| Storage: | -20 °C |
| Storage Comment: | Store between -10°C and -20°C |