# antibodies .- online.com





Go to Product page

### Datasheet for ABIN753709

## anti-ASAP1 antibody (AA 154-190) (Cy7)

| Overview              |  |  |
|-----------------------|--|--|
| Quantity:             | 100 μL   |  |
| Target:               | ASAP1  |  |
| Binding Specificity:  | AA 154-190   |  |
| Reactivity:           | Mouse, Human   |  |
| Host:                 | Rabbit   |  |
| Clonality:            | Polyclonal   |  |
| Conjugate:            | This ASAP1 antibody is conjugated to Cy7   |  |
| Application:          | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |
| Product Details       |  |  |
| Immunogen:            | KLH conjugated synthetic peptide derived from human ASAP1  |  |
| Isotype:              | IgG  |  |
| Cross-Reactivity:     | Human, Mouse   |  |
| Predicted Reactivity: | Rat,Dog,Cow,Horse,Rabbit   |  |

Purified by Protein A.

Purification:

**Target Details** 

Target: ASAP1

Alternative Name: ASAP1 (ASAP1 Products)

#### **Target Details**

| Doolyground         | Circonimo: DAD DACO AMADI DDEEL ZOLAD CENTRA Art CAD with CHO domain ANI/ report                      |
|---------------------|---|
| Background:         | Synonyms: PAP, PAG2, AMAP1, DDEF1, ZG14P, CENTB4, Arf-GAP with SH3 domain, ANK repeat                 |
|                     | and PH domain-containing protein 1, 130 kDa phosphatidylinositol 4,5-bisphosphate-dependent           |
|                     | ARF1 GTPase-activating protein, ADP-ribosylation factor-directed GTPase-activating protein 1,         |
|                     | ARF GTPase-activating protein 1, Development and differentiation-enhancing factor 1, DEF-1,           |
|                     | Differentiation-enhancing factor 1, PIP2-dependent ARF1 GAP, ASAP1, KIAA1249                          |
|                     | Background: Possesses phosphatidylinositol 4,5-bisphosphate-dependent GTPase-activating               |
|                     | protein activity for ARF1 (ADP ribosylation factor 1) and ARF5 and a lesser activity towards          |
|                     | ARF6. May coordinate membrane trafficking with cell growth or actin cytoskeleton remodeling           |
|                     | by binding to both SRC and PIP2. May function as a signal transduction protein involved in the        |
|                     | differentiation of fibroblasts into adipocytes and possibly other cell types (By similarity). Plays a |
|                     | role in ciliogenesis.   |
| Gene ID:            | 50807   |
| UniProt:            | Q9ULH1  |
| Offin Tot.          | Q50EIII   |
| Application Details |   |
| Application Notes:  | IF(IHC-P) 1:50-200  |
|                     | IF(IHC-F) 1:50-200  |
|                     | IF(ICC) 1:50-200  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 μg/μL   |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and           |
|                     | 50 % Glycerol.  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be                   |
|                     | handled by trained staff only.  |
| Storage:            | -20 °C  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                     |
| Expiry Date:        | 12 months   |