



[Go to Product page](#)

Datasheet for ABIN7538242 FFAR3 Protein

Overview

Quantity:	50 µg
Target:	FFAR3
Origin:	Human
Source:	Mammalian Cells
Protein Type:	Synthetic Nanodisc

Product Details

Purpose:	Human FFAR3 full length protein-synthetic nanodisc
Characteristics:	Unlike other membrane scaffold protein (MSP) Nanodisc on the market, our synthetic Nanodisc can be prepared directly from the cells. The polymers used during this process have a dual function. It dissolves the cell membranes, like the detergent, and uses cellular phospholipids to form Nanodisc around the membrane proteins. The target protein embedded Nanodiscs can then be purified.

Target Details

Target:	FFAR3
Alternative Name:	FFAR3 (FFAR3 Products)
Background:	G protein-coupled receptor that is activated by a major product of dietary fiber digestion, the short chain fatty acids (SCFAs), and that plays a role in the regulation of whole-body energy homeostasis and in intestinal immunity. In omnivorous mammals, the short chain fatty acids acetate, propionate and butyrate are produced primarily by the gut microbiome that metabolizes dietary fibers. SCFAs serve as a source of energy but also act as signaling

Target Details

molecules. That G protein-coupled receptor is probably coupled to the pertussis toxin-sensitive, G(i/o)-alpha family of G proteins. Its activation results in the formation of inositol 1,4,5-trisphosphate, the mobilization of intracellular calcium, the phosphorylation of the MAPK3/ERK1 and MAPK1/ERK2 kinases and the inhibition of intracellular cAMP accumulation (PubMed:12711604). Activated by SCFAs and by beta-hydroxybutyrate, a ketone body produced by the liver upon starvation, it inhibits N-type calcium channels and modulates the activity of sympathetic neurons through a signaling cascade involving the beta and gamma subunits of its coupled G protein, phospholipase C and MAP kinases. Thereby, it may regulate energy expenditure through the control of the sympathetic nervous system that controls for instance heart rate. Upon activation by SCFAs accumulating in the intestine, it may also signal to the brain via neural circuits which in turn would regulate intestinal gluconeogenesis. May also control the production of hormones involved in whole-body energy homeostasis. May for instance, regulate blood pressure through renin secretion. May also regulate secretion of the PYY peptide by enteroendocrine cells and control gut motility, intestinal transit rate, and the harvesting of energy from SCFAs produced by gut microbiota. May also indirectly regulate the production of LEP/Leptin, a hormone acting on the CNS to inhibit food intake, in response to the presence of short-chain fatty acids in the intestine. Finally, may also play a role in glucose homeostasis. Besides its role in energy homeostasis, may play a role in intestinal immunity. May mediate the activation of the inflammatory and immune response by SCFAs in the gut, regulating the rapid production of chemokines and cytokines by intestinal epithelial cells. Among SCFAs, the fatty acids containing less than 6 carbons, the most potent activators are probably propionate, butyrate and pentanoate while acetate is a poor activator (PubMed:12496283, PubMed:12711604).[UniProtKB/Swiss-Prot Function]

Molecular Weight: The human full length FFAR3 protein has a MW of 38.6kDa

UniProt: [O14843](#)

Application Details

Comment: Advantages of Synthetic Nanodiscs:

- Highly purified membrane proteins
- High solubility in aqueous solutions
- High stability
- Proteins are in a native membrane environment and remain biologically active
- No detergent and can be used for cell-based assays
- No MSP backbone proteins

Limitations of Synthetic Nanodiscs:

Application Details

- Intolerant to acids and high concentrations of divalent metal ions
-

Restrictions: For Research Use only

Handling

Format: Lyophilized

Buffer: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0).
Normally 5 % - 8 % trehalose is added as protectants before lyophilization.

Storage: -20 °C, -80 °C

Storage Comment: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).
Lyophilized proteins are shipped at ambient temperature.

Expiry Date: 12 months
