



[Go to Product page](#)

Datasheet for ABIN7538324
HCAR2 Protein

Overview

Quantity:	50 µg
Target:	HCAR2
Origin:	Human
Source:	Mammalian Cells
Protein Type:	Synthetic Nanodisc

Product Details

Purpose:	Human HCAR2 full length protein-synthetic nanodisc
Characteristics:	Unlike other membrane scaffold protein (MSP) Nanodisc on the market, our synthetic Nanodisc can be prepared directly from the cells. The polymers used during this process have a dual function. It dissolves the cell membranes, like the detergent, and uses cellular phospholipids to form Nanodisc around the membrane proteins. The target protein embedded Nanodiscs can then be purified.

Target Details

Target:	HCAR2
Alternative Name:	HCAR2 (HCAR2 Products)
Background:	Acts as a high affinity receptor for both nicotinic acid (also known as niacin) and (D)-beta-hydroxybutyrate and mediates increased adiponectin secretion and decreased lipolysis through G(i)-protein-mediated inhibition of adenylyl cyclase. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet. Mediates nicotinic acid-induced apoptosis in mature neutrophils. Receptor activation by nicotinic acid

Target Details

results in reduced cAMP levels which may affect activity of cAMP-dependent protein kinase A and phosphorylation of target proteins, leading to neutrophil apoptosis. The rank order of potency for the displacement of nicotinic acid binding is 5-methyl pyrazole-3-carboxylic acid = pyridine-3-acetic acid > acifran > 5-methyl nicotinic acid = acipimox >> nicotinuric acid = nicotinamide.[UniProtKB/Swiss-Prot Function]

Molecular Weight: The human full length HCAR2 protein has a MW of 41.9kDa

UniProt: [Q8TDS4](#)

Application Details

Comment: Advantages of Synthetic Nanodiscs:

- Highly purified membrane proteins
- High solubility in aqueous solutions
- High stability
- Proteins are in a native membrane environment and remain biologically active
- No detergent and can be used for cell-based assays
- No MSP backbone proteins

Limitations of Synthetic Nanodiscs:

- Intolerant to acids and high concentrations of divalent metal ions

Restrictions: For Research Use only

Handling

Format: Lyophilized

Buffer: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0).
Normally 5 % - 8 % trehalose is added as protectants before lyophilization.

Storage: -20 °C,-80 °C

Storage Comment: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).
Lyophilized proteins are shipped at ambient temperature.

Expiry Date: 12 months