

## Datasheet for ABIN7538324 **HCAR2 Protein**



[Go to Product page](#)

### Overview

Quantity:	50 µg
Target:	HCAR2
Origin:	Human
Source:	Mammalian Cells
Protein Type:	Synthetic Nanodisc

### Product Details

Purpose:	Human HCAR2 full length protein-synthetic nanodisc
Characteristics:	Unlike other membrane scaffold protein (MSP) Nanodisc on the market, our synthetic Nanodisc can be prepared directly from the cells. The polymers used during this process have a dual function. It dissolves the cell membranes, like the detergent, and uses cellular phospholipids to form Nanodisc around the membrane proteins. The target protein embedded Nanodiscs can then be purified.

### Target Details

Target:	HCAR2
Alternative Name:	HCAR2 ( <a href="#">HCAR2 Products</a> )
Background:	Acts as a high affinity receptor for both nicotinic acid (also known as niacin) and (D)-beta-hydroxybutyrate and mediates increased adiponectin secretion and decreased lipolysis through G(i)-protein-mediated inhibition of adenylyl cyclase. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet. Mediates nicotinic acid-induced apoptosis in mature neutrophils. Receptor activation by nicotinic acid

## Target Details

results in reduced cAMP levels which may affect activity of cAMP-dependent protein kinase A and phosphorylation of target proteins, leading to neutrophil apoptosis. The rank order of potency for the displacement of nicotinic acid binding is 5-methyl pyrazole-3-carboxylic acid = pyridine-3-acetic acid > acifran > 5-methyl nicotinic acid = acipimox >> nicotinuric acid = nicotinamide.[UniProtKB/Swiss-Prot Function]

Molecular Weight: The human full length HCAR2 protein has a MW of 41.9kDa

UniProt: [Q8TDS4](#)

## Application Details

Comment: Advantages of Synthetic Nanodiscs:

- Highly purified membrane proteins
- High solubility in aqueous solutions
- High stability
- Proteins are in a native membrane environment and remain biologically active
- No detergent and can be used for cell-based assays
- No MSP backbone proteins

Limitations of Synthetic Nanodiscs:

- Intolerant to acids and high concentrations of divalent metal ions

Restrictions: For Research Use only

## Handling

Format: Lyophilized

Buffer: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5 % - 8 % trehalose is added as protectants before lyophilization.

Storage: -20 °C,-80 °C

Storage Comment: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Expiry Date: 12 months