

Datasheet for ABIN7538457

Mu Opioid Receptor 1 Protein



Overview

Quantity	50 ug
Quantity:	50 μg
Target:	Mu Opioid Receptor 1 (OPRM1)
Origin:	Human
Source:	Mammalian Cells
Protein Type:	Synthetic Nanodisc

Product Details

Purpose:

Characteristics:	Unlike other membrane scaffold protein (MSP) Nanodisc on the market, our synthetic Nanodisc
	can be prepared directly from the cells. The polymers used during this process have a dual
	function. It dissolves the cell membranes, like the detergent, and uses cellular phospholipids to
	form Nanodisc around the membrane proteins. The target protein embedded Nanodiscs can
	then be purified.

Human OPRM full length protein-synthetic nanodisc

Target Details

Target:	Mu Opioid Receptor 1 (OPRM1)
Alternative Name:	OPRM (OPRM1 Products)
Background:	This gene encodes one of at least three opioid receptors in humans, the mu opioid receptor
	(MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic
	agents such as beta-endorphin and enkephalins. The MOR also has an important role in
	dependence to other drugs of abuse, such as nicotine, cocaine, and alcohol via its modulation
	of the dopamine system. The NM_001008503.2:c.118A>G allele has been associated with

	opioid and alcohol addiction and variations in pain sensitivity but evidence for it having a causal role is conflicting. Multiple transcript variants encoding different isoforms have been found for this gene. Though the canonical MOR belongs to the superfamily of 7-transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only 6 transmembrane domains. [provided by RefSeq, Oct 2013]
Molecular Weight:	The human full length OPRM protein has a MW of 44.8kDa
UniProt:	P35372
Pathways:	cAMP Metabolic Process, Synaptic Membrane

Application Details

Comment:	Advantages of

Advantages of Synthetic Nanodiscs:

- · Highly purified membrane proteins
- · High solubility in aqueous solutions
- High stability
- · Proteins are in a native membrane environment and remain biologically active
- · No detergent and can be used for cell-based assays
- No MSP backbone proteins

Limitations of Synthetic Nanodiscs:

• Intolerant to acids and high concentrations of divalent metal ions

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Buffer:	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Expiry Date:	12 months