

Datasheet for ABIN7538512 **RBM24 Protein**



Overview

Quantity:	50 µg
Target:	RBM24
Origin:	Human
Source:	Mammalian Cells
Protein Type:	Synthetic Nanodisc

Product Details

Purpose:	Human MTR1A full length protein-synthetic nanodisc
Characteristics:	Unlike other membrane scaffold protein (MSP) Nanodisc on the market, our synthetic Nanodisc can be prepared directly from the cells. The polymers used during this process have a dual
	function. It dissolves the cell membranes, like the detergent, and uses cellular phospholipids to
	form Nanodisc around the membrane proteins. The target protein embedded Nanodiscs can
	then be purified.

Target Details

Target:	RBM24
Alternative Name:	MTR1A (RBM24 Products)
Background:	This gene encodes one of two high affinity forms of a receptor for melatonin, the primary
	hormone secreted by the pineal gland. This receptor is a G-protein coupled, 7-transmembrane
	receptor that is responsible for melatonin effects on mammalian circadian rhythm and
	reproductive alterations affected by day length. The receptor is an integral membrane protein
	that is readily detectable and localized to two specific regions of the brain. The hypothalamic

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7538512 | 07/24/2024 | Copyright antibodies-online. All rights reserved.

	suprachiasmatic nucleus appears to be involved in circadian rhythm while the hypophysial pars
	tuberalis may be responsible for the reproductive effects of melatonin. [provided by RefSeq, Jul
	2008]
Molecular Weight:	The human full length MTR1A protein has a MW of 39.4kDa
UniProt:	P48039
Pathways:	Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development

Application Details

Comment:	Advantages of Synthetic Nanodiscs:
	Highly purified membrane proteins
	High solubility in aqueous solutions
	High stability
	Proteins are in a native membrane environment and remain biologically active
	No detergent and can be used for cell-based assays
	No MSP backbone proteins
	Limitations of Synthetic Nanodiscs:
	Intolerant to acids and high concentrations of divalent metal ions
Restrictions:	For Research Use only
Handling	
Handling Format:	Lyophilized
_	Lyophilized Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0).
Format:	
Format:	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0).
Format: Buffer:	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
Format: Buffer: Storage:	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5 % - 8 % trehalose is added as protectants before lyophilization. -20 °C,-80 °C
Format: Buffer: Storage:	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0).Normally 5 % - 8 % trehalose is added as protectants before lyophilization20 °C,-80 °CStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for