

Datasheet for ABIN7538871

anti-Cytokeratin 5 antibody (C-Term)





Overview


Quantity:	1 mL
Target:	Cytokeratin 5 (KRT5)
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Cytokeratin 5 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Heat-induced Epitope Retrieval (HIER)
Product Details	
Product Details Purpose:	Liquid Mouse Monoclonal Antibody Cytokeratin 5
	Liquid Mouse Monoclonal Antibody Cytokeratin 5 Prokaryotic recombinant fusion protein corresponding to a 103 amino acid portion of the C-terminal region of the human cytokeratin 5 molecule.
Purpose:	Prokaryotic recombinant fusion protein corresponding to a 103 amino acid portion of the C-
Purpose: Immunogen:	Prokaryotic recombinant fusion protein corresponding to a 103 amino acid portion of the C-terminal region of the human cytokeratin 5 molecule.
Purpose: Immunogen: Clone:	Prokaryotic recombinant fusion protein corresponding to a 103 amino acid portion of the C-terminal region of the human cytokeratin 5 molecule. XM26
Purpose: Immunogen: Clone: Isotype:	Prokaryotic recombinant fusion protein corresponding to a 103 amino acid portion of the C-terminal region of the human cytokeratin 5 molecule. XM26 IgG1, kappa

Target Details

Expiry Date:

12 months

Target Details	
Target:	Cytokeratin 5 (KRT5)
Alternative Name:	Cytokeratin 5 (KRT5 Products)
Application Details	
Application Notes:	Immunohistochemistry on paraffin sections. Heat Induced Epitope Retrieval (HIER): Epitope Retrieval Solution pH 6. Suggested dilution: 1:100 for 30 minutes at 25 °C. This is provided as a guide and users should determine their own optimal working dilutions. Immunohistochemical (IHC) staining techniques allow for the visualization of antigens via the sequential application of a specific antibody to the antigen (primary antibody), a secondary antibody to the primary antibody and an enzyme complex with a chromogenic substrate with interposed washing steps. The enzymatic activation of the chromogen results in a visible reaction product at the antigen site. The specimen may then be counterstained and coverslipped. Results are interpreted using a light microscope.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	21.0 mg/L
Buffer:	This product is a liquid tissue culture supernatant containing sodium azide as a preservative.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C. Do not freeze. Return to 2-8 °C immediately after use. Do not use after expiration date indicated on the vial label. Storage conditions other than those specified above must be verified by the user.



Immunohistochemistry

Image 1. Human breast, ductal carcinoma in situ: immunohistochemical staining for Cytokeratin 5. Note the intense staining of myoepithelial cells. Cytokeratin 5: clone XM26