

Datasheet for ABIN7538883

anti-MUC1 antibody





Overview

Quantity:	1 mL
Target:	MUC1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This MUC1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),
	Heat-induced Epitope Retrieval (HIER)

Product Details

Purpose:	Liquid Mouse Monoclonal Antibody Epithelial Membrane Antigen
Immunogen:	Human milk fat globule membrane.
Clone:	GP1-4
Isotype:	lgG1
Specificity:	Human epithelial membrane antigen.
Characteristics:	This product is intended for the qualitative identification by light microscopy of Epithelial Membrane Antigen molecules in paraffin sections.
Purification:	Tissue culture supernatant

Target Details

Target:	MUC1
Alternative Name:	Epithelial Membrane Antigen (MUC1 Products)
Pathways:	Negative Regulation of intrinsic apoptotic Signaling

Application Details


Immunohistochemistry on paraffin sections. Heat Induced Epitope Retrieval (HIER): Epitope Retrieval Solution pH 6. Suggested dilution: 1:200 for 30 minutes at 25 °C. This is provided as a guide and users should determine their own optimal working dilutions. Immunohistochemical (IHC) staining techniques allow for the visualization of antigens via the sequential application of a specific antibody to the antigen (primary antibody), a secondary antibody to the primary antibody and an enzyme complex with a chromogenic substrate with interposed washing steps. The enzymatic activation of the chromogen results in a visible reaction product at the antigen site. The specimen may then be counterstained and coverslipped. Results are interpreted using a light microscope.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	11 mg/L
Buffer:	This product is a liquid tissue culture supernatant containing sodium azide as a preservative.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C. Do not freeze. Return to 2-8 °C immediately after use. Do not use after expiration date indicated on the vial label. Storage conditions other than those specified above must be verified by the user.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Human breast cancer: immunohistochemical staining for Epithelial Membrane Antigen. Note membrane staining of neoplastic cells. Epithelial Membrane Antigen: clone GP1.4