

Datasheet for ABIN753898

anti-Filamin A antibody (AA 1851-1960)



_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Overview		
Quantity:	100 μL	
Target:	Filamin A (FLNA)	
Binding Specificity:	AA 1851-1960	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Filamin A antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human Filamin A	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	Filamin A (FLNA)	

Target Details

Alternative Name:	Filamin A (FLNA Products)	
Background:	Synonyms: ABP 280, ABP 280 like protein, ABP-280, ABP280A, ABPA, Actin binding like protein,	
	Actin binding protein 280, Actin-binding protein 280, Alpha filamin, Alpha-filamin, APBX, cb967,	
	Dilp2, Endothelial actin binding protein, Endothelial actin-binding protein, Filamin 1, Filamin A	
	alpha actin binding protein 280, Filamin A, Filamin-1, Filamin-A, FLN, FLN-A, FLN1, FLNA,	
	FLNA_HUMAN, FMD, MNS, NHBP, Non muscle filamin, Non-muscle filamin, OPD, OPD1, OPD2.	
	Background: Caldesmon, Filamin 1, Nebulin and Villin are differentially expressed and regulated	
	Actin binding proteins. Both muscular (CDh) and non-muscular (CDI) forms of Caldesmon have	
	been identified and each has been shown to bind to Actin as well as to calmodulin and Myosin.	
	CDh is expressed predominantly on thin filaments in smooth muscle, whereas CDI is widely	
	expressed in non-muscle tissues and cells. Filamin 1, which is ubiquitously expressed and	
	exists as a homodimer, functions to crosslink Actin to filaments. Nebulin is a large filamentous	
	protein specific to muscle tissue that may function as a ruler for filament length. Several	
	isoforms of Nebulin are produced by alternative exon usage. Villin is Ca2+-regulated and is the	
	major structural component of the brush border of absorptive cells.	
Gene ID:	2316	
Pathways:	TCR Signaling, Maintenance of Protein Location	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	

Handling

Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	