

Datasheet for ABIN7540532 anti-eEF1A1 antibody (N-Term)



| _ | | | | | |
|---|---|---|----|----|---|
| | W | 0 | rv | 10 | W |

| Quantity: | 100 μg | |
|-----------------------------|---|--|
| Target: | eEF1A1 (EEF1A1) | |
| Binding Specificity: | N-Term | |
| Reactivity: | Human | |
| Host: | Chicken | |
| Clonality: | Polyclonal | |
| Conjugate: | This eEF1A1 antibody is un-conjugated | |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC) | |
| Product Details | | |
| Purpose: | EEF1A1 Antibody | |
| Immunogen: | Anti-EEF1A1 antibody was prepared from eggs of chickens laid after repeated immunizations with a synthetic peptide corresponding to a N-terminal portion of human EEF1A1 conjugated to Keyhole Limpet Hemocyanin (KLH). | |
| Isotype: | IgG | |
| Cross-Reactivity (Details): | This affinity purified antibody is directed against human EEF1A1. | |
| Purification: | This product is an IgY fraction antibody purified from monospecific chicken egg yolks by a multi-step process which includes selective precipitation and salt fractionation followed by extensive dialysis against the buffer stated above. | |
| Sterility: | Sterile filtered | |

Target Details

| Target: | eEF1A1 (EEF1A1) | | |
|---------------------|--|--|--|
| Alternative Name: | EEF1A1 (EEF1A1 Products) | | |
| Background: | Chicken Anti-Eukaryotic Translation Elongation Factor 1 Alpha 1, Leukocyte Receptor Cluster | | |
| | (LRC) Member 7, Leukocyte Receptor Cluster Member 7, Eukaryotic Elongation Factor 1 A-1, | | |
| | Elongation Factor 1-Alpha 1, Elongation Factor Tu, EF-1-Alpha-1, EEF1A-1, EEF1A, LENG7, EF- | | |
| | Tu, EF1A, Eukaryotic Translation Elongation Factor 1 Alpha 1-Like 14, Glucocorticoid Receptor | | |
| | AF-1 Specific Elongation Factor, Translation Elongation Factor 1 Alpha 1-Like 14, Epididymis | | |
| | Secretory Sperm Binding Protein, Elongation Factor 1 Alpha Subunit, Prostate Tumor-Inducing | | |
| | Protein 1, Cervical Cancer Suppressor 3, CTCL Tumor Antigen, EF1a-Like Protein, HNGC:16303 | | |
| | GRAF-1EF, CCS-3, EE1A1, EEF-1, CCS3, PTI1,EEF1A1 (Eukaryotic Translation Elongation Factor | | |
| | 1 Alpha 1) protein promotes the GTP-dependent binding of aminoacyl-tRNA to the A-site of | | |
| | ribosomes during protein biosynthesis. With PARP1 and TXK, forms a complex that acts as a T | | |
| | helper 1 (Th1) cell-specific transcription factor and binds the promoter of IFN-gamma to | | |
| | directly regulate its transcription, and is thus involved importantly in Th1 cytokine production. | | |
| | This isoform (alpha 1) is expressed in brain, placenta, lung, liver, kidney, and pancreas, and the | | |
| | other isoform (alpha 2) is expressed in brain, heart and skeletal muscle. This isoform is | | |
| | identified as an autoantigen in 66 % of patients with Felty syndrome. Anti-EEF1A1 is useful for | | |
| | researchers interested in Eukaryotic Translation Elongation Factor 1 Alpha-1-Like 14, Cervical | | |
| | Cancer, and the Innate Immune system. | | |
| Gene ID: | 1915 | | |
| NCBI Accession: | NP_001393 | | |
| UniProt: | P68104 | | |
| Application Details | | | |
| Application Notes: | ELISA_Dilution: 1:10,000 - 1:50,000 | | |
| | Immunohistochemistry_Dilution: 1:100 | | |
| | Western_Blot_Dilution: 5 µg/mL | | |
| Comment: | Anti-EEF1A1 Antibody has been tested in Western Blot and IHC. Expect a band at ~48-50 kDa | | |
| | in western blot using appropriate tissues and lysates. Positive control used: MCF-7 nuclear | | |
| | extract lysate in western blot and human kidney tissue in Immunohistochemistry. | | |
| Restrictions: | For Research Use only | | |
| | | | |

Handling

| Format: | Liquid | |
|--------------------|---|--|
| Buffer: | Buffer: 0.002 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.0 Stabilizer: None Preservative: 0.01 % (w/v) Sodium Azide | |
| Preservative: | Sodium azide | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | 4 °C,-20 °C | |
| Storage Comment: | Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. | |
| Expiry Date: | 12 months | |