

# Datasheet for ABIN7540567 anti-EEF1A2 antibody (C-Term)



Quantity:	100 μg	
Target:	EEF1A2	
Binding Specificity:	C-Term	
Reactivity:	Human	
Host:	Goat	
Clonality:	Polyclonal	
Conjugate:	This EEF1A2 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Fluorescence Microscopy (FM)	

### **Product Details**

Overview

Purpose:	EEF1A2 Antibody
Immunogen:	Anti-EEF1A2 antibody was prepared from whole goat serum produced by repeated immunizations with a synthetic peptide corresponding to a near C-terminal portion of human EEF1A2 conjugated to Keyhole Limpet Hemocyanin (KLH).
Isotype:	IgG
Cross-Reactivity (Details):	This affinity purified antibody is directed against human EEF1A2.
Purification:	The product was affinity purified from monospecific antiserum by immunoaffinity purification.
Sterility:	Sterile filtered

# Target Details

Target:	EEF1A2
Alternative Name:	EEF1A2 (EEF1A2 Products)
Background:	Goat Anti-EEF1A2 Antibody, Goat Anti-Eukaryotic Translation Elongation Factor 1 Alpha 2
	Antibody, Eukaryotic Elongation Factor 1 A-2, EF-1-Alpha-2, Statin-S1, EEF1AL, STN 3,
	Elongation Factor 1-Alpha 2, Statin-Like, EEF1A-2, EIEE33, Statin, MRD38, EF1A, STNL,
	HS1,EEF1A2 (Eukaryotic Translation Elongation Factor 1 Alpha 2) protein promotes the GTP-
	dependent binding of aminoacyl-tRNA to the A-site of ribosomes during protein biosynthesis.
	EEF1A2 encodes an isoform of the alpha subunit of the elongation factor-1 complex, which is
	responsible for the enzymatic delivery of aminoacyl tRNAs to the ribosome. This isoform (alpha
	2) is expressed in brain, heart and skeletal muscle, and the other isoform (alpha 1) is expressed
	in brain, placenta, lung, liver, kidney, and pancreas. EEF1A2 may be associated with the
	development of ovarian cancer, along with Epileptic Encephalopathy, Early Infantile, 33 and
	Mental Retardation, Autosomal Dominant 38. Anti-EEF1A2 Antibody is useful for researchers
	interested in cancer research, translation factors, GTP binding, and GTPase activities.
Gene ID:	1917
NCBI Accession:	NP_001949
UniProt:	Q05639
Application Details	
Application Notes:	ELISA_Dilution: 1:10,000-1:50,000
	Immunohistochemistry_Dilution: 1:200
	IF_Microscopy_Dilution: 15 μg/mL
	Western_Blot_Dilution: 1:1000
Comment:	Anti-EEF1A2 Antibody has been tested in WB, IF, and IHC. Expect a band at ~50.5 kDa in
	western blot using appropriate lysates. Positive control used: human skeletal muscle lysate,
	MCF-7 nuclear extract lysate in WB, MCF-7 cells in IF, and human skeletal muscle tissue in IHC.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
	Stabilizer: None

# Handling

	Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiry Date:	12 months