# antibodies .- online.com







## anti-PPAP2B antibody (AA 241-311) (Cy5.5)



O	:	
Ove	r \/ I	$\Theta \backslash \Lambda /$
$\circ$	1 V I	$\sim$ v v

Quantity:	100 μL
Target:	PPAP2B
Binding Specificity:	AA 241-311
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PPAP2B antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PAP2B
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

#### **Target Details**

Target:	PPAP2B
Alternative Name:	PAP2B (PPAP2B Products)

### Target Details

9	
Background:	Synonyms: LPP3, VCIP, Dri42, PAP2B, Lipid phosphate phosphohydrolase 3, PAP2-beta,
	Phosphatidate phosphohydrolase type 2b, Phosphatidic acid phosphatase 2b, PAP-2b, Vascular
	endothelial growth factor and type I collagen-inducible protein, PPAP2B
	Background: Catalyzes the conversion of phosphatidic acid (PA) to diacylglycerol (DG). In
	addition it hydrolyzes lysophosphatidic acid (LPA), ceramide-1-phosphate (C-1-P) and
	sphingosine-1-phosphate (S-1-P). The relative catalytic efficiency is LPA = PA > C-1-P > S-1-P.
	May be involved in cell adhesion and in cell-cell interactions.
Gene ID:	8613
UniProt:	014495
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months