

Datasheet for ABIN7543100

Recombinant anti-MOG antibody (AA 1-125)



_				
()	1//	rv	IO	Λ/
()	VC	. I V	1	v v

Quantity:	200 μg
Target:	MOG
Binding Specificity:	AA 1-125
Reactivity:	Rat
Host:	Rat
Antibody Type:	Recombinant Antibody
Clonality:	Chimeric
Conjugate:	This MOG antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA, Blocking Peptide (BP)

Product Details

Purpose:	Anti-MOG [M3-8], Rat IgG1, kappa	
Immunogen:	Callithrix jacchus marmosets were used, and experimental allergic encephalomyelitis was	
	induced by the injection of rat MOG (aa1-125) into the marmosettes. The rMOG was expressed	
	in E. coli and purified to homogeneity. The animals were killed 4-70 days after the onset of	
	symptoms of EAE. Bone marrow and spleen cells were obtained from an immunized C. jacchus	
	the RNA extracted with Trizol reagent and rtPCR used to generate cloning inserts containing	
	Fab portions of IgGk. Phage display was then used to select for the MOG-reactive Fab	
	fragments using the pCOMB3H phage display vector, and binding confirmed using an ELISA.	
Clone:	M3-8	
Isotype:	lgG1 kappa	

Product Details

Restrictions:	For Research Use only	
Comment:	This chimeric rat antibody was made using the variable domain sequences of the original Marmoset Fab format, for improved compatibility with existing reagents, assays and techniques.	
Application Notes:	This antibody is part of a family of antibodies including clones M26, M38, M45, M3-8, M3-24, M3-31. This antibody has been proposed for the diagnosis and prognosis of multiple sclerosis (MS) or experimental allergic encephalomyelitis (EAE) which is a disease model for MS. This is achieved by using competition assays to determine if there are autoantibodies present in an individual which recognise structural epitopes on MOG which have been shown to be associated with the progression of MS. This antibody has been used in ELISAs and competition assays to characterise its epitope (see specificity statement) and determine whether the similar epitopes are recognised in marmosets with EAE and humans with MS. It was originally generated and tested as a Fab (von Budingen et al, 2002). Whilst this has been shown to not bind to hMOG, it has been shown to compete with IgG from MS patients for binding to rMOG (Lalive et al, 2006). Analysis has also been done on the amino acid sequence of this antibody (von Budingen et al, 2006).	
Application Details		
UniProt:	Q63348	
Background:	Myelin oligodendrocyte glycoprotein, rMOG	
Alternative Name:	MOG (MOG Products)	
Target:	MOG	
Target Details		
Purification:	Protein A affinity purified	
Characteristics:	Original Species of Ab: Common Marmoset Original Format of Ab: Fab	
	immunofluorescence. The epitope recognised is a structural epitope, shown by the lack of recognition of linear MOG peptides. Has also been shown to compete with anti-MOG Abs from three patients with MS. Does not bind to MOG expressed in CHO cells.	
Specificity:	Recognises an epitope which is distinct from M26, M38, M45, M3-24 and M3-31. Is able to displace native anti-MOG Abs from C. jacchus serum. Has been shown to bind in situ by	

Handling

Concentration:	1 mg/mL	
Buffer:	PBS with 0.02 % Proclin 300.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.	