antibodies -online.com





anti-Neurofibromin 1 antibody (AA 2701-2839) (Biotin)



()	1 /	0	rv	/ 1 /	71	Α.
	1//	\vdash	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL	
Target:	Neurofibromin 1 (NF1)	
Binding Specificity:	AA 2701-2839	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Neurofibromin 1 antibody is conjugated to Biotin	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Neurofibromin 1	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Rat,Dog,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	Neurofibromin 1 (NF1)
Alternative Name:	NF1 (NF1 Products)

Target Details

Background:	Synonyms: Neurofibromin 1, DKFZp686J1293, FLJ21220, Neurofibromatosis Noonan
	syndrome, Neurofibromatosis related protein NF 1, Neurofibromatosis related protein NF1,
	neurofibromatosis type I, Neurofibromatosis-related protein NF-1, Neurofibromin 1,
	Neurofibromin truncated, Neurofibromin1, NF 1, NF, NF1, NF1_HUMAN, NFNS, Type 1
	Neurofibromatosis, von Recklinghausen disease neurofibromin, von Recklinghausen disease
	related protein VRNF, VRNF, WATS, Watson disease related protein WSS, Watson syndrome, WSS.
	Background: This gene product appears to function as a negative regulator of the ras signal
	transduction pathway. Mutations in this gene have been linked to neurofibromatosis type 1,
	juvenile myelomonocytic leukemia and Watson syndrome. The mRNA for this gene is subject to
	RNA editing (CGA>UGA->Arg1306Term) resulting in premature translation termination.
	Alternatively spliced transcript variants encoding different isoforms have also been described
	for this gene. [provided by RefSeq, Jul 2008].
Gene ID:	4763
UniProt:	P21359
Pathways:	cAMP Metabolic Process, Regulation of long-term Neuronal Synaptic Plasticity
Application Details	
Application Notes:	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
	This was dust as attains Dus Oline a DOLONNOLIO AND LIAZADDOLIO OLIDOTANOE subjek abaseld be
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
Precaution of Use:	handled by trained staff only.

Handling

Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	