

Datasheet for ABIN754618 anti-PAK6 antibody (AA 511-610)



Overview

Quantity:	100 μL
Target:	PAK6
Binding Specificity:	AA 511-610
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PAK6 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PAK6
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

rardet. PAN	AK6					

Target Details

Alternative Name:	PAK6 (PAK6 Products)				
Background:	Synonyms: CDKN1A activated kinase 6, p21 activated protein kinase 6, p21 protein Cdc42/Rac-				
	activated kinase 6, p21CDKN1A activated kinase 6, p21-ACTIVATED KINASE 6, p21activated				
	kinase 6, PAK 5, PAK 6, PAK5, Serine threonine protein kinase PAK 6, Serine/threonine protein				
	kinase PAK 6, Serine/threonine protein kinase PAK6, EC 2.7.1.37, PAK-5, PAK-6, Pak6,				
	PAK6_HUMAN.				
	Background: This gene encodes a protein that shares a high degree of sequence similarity with				
	p21-activated kinase (PAK) family members. The proteins of this family are Rac/Cdc42-				
	associated Ste20-like Ser/Thr protein kinases, characterized by a highly conserved amino-				
	terminal Cdc42/Rac interactive binding (CRIB) domain and a carboxyl-terminal kinase domain.				
	PAK kinases are implicated in the regulation of a number of cellular processes, including				
	cytoskeleton rearrangement, apoptosis and the MAP kinase signaling pathway. The protein				
	encoded by this gene was found to interact with androgen receptor (AR), which is a steroid				
	hormone-dependent transcription factor that is important for male sexual differentiation and				
	development. The p21-activated protein kinase 6 gene was found to be highly expressed in				
	testis and prostate tissues and the encoded protein was shown to cotranslocate into the				
	nucleus with AR in response to androgen.				
Gene ID:	56924				
Application Details					
Application Notes:	ELISA 1:500-1000				
	IHC-P 1:200-400				
	IHC-F 1:100-500				
	IF(IHC-P) 1:50-200				
	IF(IHC-F) 1:50-200				
	IF(ICC) 1:50-200				
Restrictions:	For Research Use only				
Handling					
Format:	Liquid				
Concentration:	1 μg/μL				
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.				
Preservative:	ProClin				

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months