

Datasheet for ABIN7546217 USP39 Protein (AA 1-565) (His tag)



Overview

Quantity:	1 mg
Target:	USP39
Protein Characteristics:	AA 1-565
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This USP39 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Purpose:	Custom-made recombinat USP39 Protein expressed in mammalien cells.
Sequence:	MSGRSKRESR GSTRGKRESE SRGSSGRVKR ERDREREPEA ASSRGSPVRV KREFEPASAR
	EAPASVVPFV RVKREREVDE DSEPEREVRA KNGRVDSEDR RSRHCPYLDT INRSVLDFDF
	EKLCSISLSH INAYACLVCG KYFQGRGLKS HAYIHSVQFS HHVFLNLHTL KFYCLPDNYE
	IIDSSLEDIT YVLKPTFTKQ QIANLDKQAK LSRAYDGTTY LPGIVGLNNI KANDYANAVL
	QALSNVPPLR NYFLEEDNYK NIKRPPGDIM FLLVQRFGEL MRKLWNPRNF KAHVSPHEML
	QAVVLCSKKT FQITKQGDGV DFLSWFLNAL HSALGGTKKK KKTIVTDVFQ GSMRIFTKKL
	PHPDLPAEEK EQLLHNDEYQ ETMVESTFMY LTLDLPTAPL YKDEKEQLII PQVPLFNILA
	KFNGITEKEY KTYKENFLKR FQLTKLPPYL IFCIKRFTKN NFFVEKNPTI VNFPITNVDL
	REYLSEEVQA VHKNTTYDLI ANIVHDGKPS EGSYRIHVLH HGTGKWYELQ DLQVTDILPQ
	MITLSEAYIQ IWKRRDNDET NQQGA Sequence without tag. The proposed Purification-Tag is
	based on experiences with the expression system, a different complexity of the protein

could make another tag necessary. In case you have a special request, please contact us. Characteristics: Key Benefits: Made to order protein - from design to production - by highly experienced protein experts. Protein expressed in mammalien cells and purified in one-step affinity chromatography · The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins. • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein. If you are not interested in a full length protein, please contact us for individual protein fragments. The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. > 90 % as determined by Bis-Tris Page, Western Blot Purity: custom-made Grade: **Target Details** USP39 Target: Alternative Name: USP39 (USP39 Products) Background: Ubiquitin carboxyl-terminal hydrolase 39 (EC 3.4.19.12) (SAD1 homolog) (U4/U6.U5 tri-snRNPassociated 65 kDa protein), FUNCTION: Deubiquitinating enzyme that plays a role in many cellular processes including cellular antiviral response, epithelial morphogenesis, DNA repair or B-cell development (PubMed:33127822, PubMed:34614178). Plays a role in pre-mRNA splicing as a component of the U4/U6-U5 tri-snRNP, one of the building blocks of the precatalytic spliceosome (PubMed:11350945, PubMed:26912367). Specifically regulates immunoglobulin gene rearrangement in a spliceosome-dependent manner, which involves modulating

chromatin interactions at the Igh locus and therefore plays an essential role in B-cell

development (By similarity). Regulates AURKB mRNA levels, and thereby plays a role in

cytokinesis and in the spindle checkpoint (PubMed:18728397). Regulates apoptosis and G2/M

(PubMed:30771428). Plays also an important role in DNA repair by controlling the recruitment

cell cycle checkpoint in response to DNA damage by deubiquitinating and stabilizing CHK2

Expiry Date:

12 months

of XRCC4/LIG4 to DNA double-strand breaks for non-homologous end-joining repair
(PubMed:34614178). Participates in antiviral activity by affecting the type I IFN signaling by
stabilizing STAT1 and decreasing its 'Lys-6'-linked ubiquitination (PubMed:33127822).
Contributes to non-canonical Wnt signaling during epidermal differentiation (By similarity). Acts
as a negative regulator NF-kappa-B activation through deubiquitination of 'Lys-48'-linked
ubiquitination of NFKBIA (PubMed:36651806). {ECO:0000250 UniProtKB:Q3TIX9,
ECO:0000269 PubMed:11350945, ECO:0000269 PubMed:18728397,
ECO:0000269 PubMed:26912367, ECO:0000269 PubMed:30771428,
ECO:0000269 PubMed:33127822, ECO:0000269 PubMed:34614178,
ECO:0000269 PubMed:36651806}.
65.4 kDa
Q53GS9
Ribonucleoprotein Complex Subunit Organization
In addition to the applications listed above we expect the protein to work for functional studies
as well. As the protein has not been tested for functional studies yet we cannot offer a
guarantee though.
For Research Use only
Liquid
The buffer composition is at the discretion of the manufacturer.
Avoid repeated freeze-thaw cycles.
-80 °C
Store at -80°C.