

Datasheet for ABIN754624 anti-PAK6 antibody (AA 511-610) (Cy7)



Overview	
Quantity:	100 μL
Target:	PAK6
Binding Specificity:	AA 511-610
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PAK6 antibody is conjugated to Cy7
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Product Details Immunogen:	KLH conjugated synthetic peptide derived from human PAK6
	KLH conjugated synthetic peptide derived from human PAK6
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Mouse
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	IgG Mouse Human,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	IgG Mouse Human,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit

Target Details

Background:

Synonyms: CDKN1A activated kinase 6, p21 activated protein kinase 6, p21 protein Cdc42/Racactivated kinase 6, p21CDKN1A activated kinase 6, p21-ACTIVATED KINASE 6, p21activated kinase 6, PAK 5, PAK 6, PAK5, Serine threonine protein kinase PAK 6, Serine/threonine protein kinase PAK 6, Serine/threonine protein kinase PAK6, EC 2.7.1.37, PAK-5, PAK-6, Pak6, PAK6_HUMAN.

Background: This gene encodes a protein that shares a high degree of sequence similarity with p21-activated kinase (PAK) family members. The proteins of this family are Rac/Cdc42-associated Ste20-like Ser/Thr protein kinases, characterized by a highly conserved aminoterminal Cdc42/Rac interactive binding (CRIB) domain and a carboxyl-terminal kinase domain. PAK kinases are implicated in the regulation of a number of cellular processes, including cytoskeleton rearrangement, apoptosis and the MAP kinase signaling pathway. The protein encoded by this gene was found to interact with androgen receptor (AR), which is a steroid hormone-dependent transcription factor that is important for male sexual differentiation and development. The p21-activated protein kinase 6 gene was found to be highly expressed in testis and prostate tissues and the encoded protein was shown to cotranslocate into the nucleus with AR in response to androgen.

Gene ID:

56924

Application Details

An	plication	n Notes:
, ,	phoduo	I I VOLCO.

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 $\%$ BSA, 0.03 $\%$ Proclin300 and 50 $\%$ Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months