

Datasheet for ABIN7546735

CYP2S1 Protein (AA 1-504) (His tag)



()	V		rV	ĺ	9	V	V
'	\mathcal{I}	٧V	<u> </u>	v	1	$\overline{}$	٧	٧

Quantity:	1 mg
Target:	CYP2S1
Protein Characteristics:	AA 1-504
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CYP2S1 protein is labelled with His tag.

Product Details

Purpose:	Custom-made recombinant CYP2S1 Protein expressed in mammalian cells.
Sequence:	MEATGTWALL LALALLLLLT LALSGTRARG HLPPGPTPLP LLGNLLQLRP GALYSGLMRL
	SKKYGPVFTI YLGPWRPVVV LVGQEAVREA LGGQAEEFSG RGTVAMLEGT FDGHGVFFSN
	GERWRQLRKF TMLALRDLGM GKREGEELIQ AEARCLVETF QGTEGRPFDP SLLLAQATSN
	VVCSLLFGLR FSYEDKEFQA VVRAAGGTLL GVSSQGGQTY EMFSWFLRPL PGPHKQLLHH
	VSTLAAFTVR QVQQHQGNLD ASGPARDLVD AFLLKMAQEE QNPGTEFTNK NMLMTVIYLL
	FAGTMTVSTT VGYTLLLLMK YPHVQKWVRE ELNRELGAGQ APSLGDRTRL PYTDAVLHEA
	QRLLALVPMG IPRTLMRTTR FRGYTLPQGT EVFPLLGSIL HDPNIFKHPE EFNPDRFLDA
	DGRFRKHEAF LPFSLGKRVC LGEGLAKAEL FLFFTTILQA FSLESPCPPD TLSLKPTVSG
	LFNIPPAFQL QVRPTDLHST TQTR Sequence without tag. The proposed Purification-Tag is
	based on experiences with the expression system, a different complexity of the protein
	could make another tag necessary. In case you have a special request, please contact us.
Specificity:	If you are looking for a specific domain and are interested in a partial protein or a different

Product Details

	isoform, please contact us regarding an individual offer.
Characteristics:	Key Benefits:
	Made to order protein - from design to production - by highly experienced protein experts.
	Protein expressed in mammalian cells and purified in one-step affinity chromatography
	The optimized expression system ensures reliability for intracellular, secreted and
	transmembrane proteins.
	State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made-to-order protein and will be made for the first time for your order. Our
	experts in the lab try to ensure that you receive soluble protein.
	If you are not interested in a full length protein, please contact us for individual protein
	fragments.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom
	made proteins from other companies is that there is no financial obligation in case the protein
	cannot be expressed or purified.
Purity:	> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)
Grade:	custom-made
Target Details	
Target:	CYP2S1
Alternative Name:	CYP2S1 (CYP2S1 Products)
Background:	
Backgrouna.	Cytochrome P450 2S1 (EC 1.14.14) (CYPIIS1) (Hydroperoxy icosatetraenoate dehydratase)
Buckground.	Cytochrome P450 2S1 (EC 1.14.14) (CYPIIS1) (Hydroperoxy icosatetraenoate dehydratase) (EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5),FUNCTION: A cytochrome P450
Buonground.	
Buonground.	(EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5),FUNCTION: A cytochrome P450
Buonground.	(EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5), FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:12711469, PubMed:21068195). In epidermis, may contribute to the oxidative metabolism of all-trans-
Buonground.	(EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5), FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:12711469, PubMed:21068195). In epidermis, may contribute to the oxidative metabolism of all-transretinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a
Buonground.	(EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5), FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:12711469, PubMed:21068195). In epidermis, may contribute to the oxidative metabolism of all-transretinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by
Buonground.	(EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5), FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:12711469, PubMed:21068195). In epidermis, may contribute to the oxidative metabolism of all-transretinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPHhemoprotein reductase)
Buonground.	(EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5), FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:12711469, PubMed:21068195). In epidermis, may contribute to the oxidative metabolism of all-transretinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPHhemoprotein reductase) (PubMed:12711469). Additionally, displays peroxidase and isomerase activities toward various
Buonground.	(EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5),FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:12711469, PubMed:21068195). In epidermis, may contribute to the oxidative metabolism of all-transretinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPHhemoprotein reductase) (PubMed:12711469). Additionally, displays peroxidase and isomerase activities toward various oxygenated eicosanoids such as prostaglandin H2 (PGH2) and hydroperoxyeicosatetraenoates
Buonground.	(EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5),FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:12711469, PubMed:21068195). In epidermis, may contribute to the oxidative metabolism of all-transretinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPHhemoprotein reductase) (PubMed:12711469). Additionally, displays peroxidase and isomerase activities toward various oxygenated eicosanoids such as prostaglandin H2 (PGH2) and hydroperoxyeicosatetraenoates (HPETEs) (PubMed:21068195). Independently of cytochrome P450 reductase, NADPH, and O2,
Background.	(EC 4.2.1.152) (Thromboxane-A synthase) (EC 5.3.99.5),FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:12711469, PubMed:21068195). In epidermis, may contribute to the oxidative metabolism of all-transretinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPHhemoprotein reductase) (PubMed:12711469). Additionally, displays peroxidase and isomerase activities toward various oxygenated eicosanoids such as prostaglandin H2 (PGH2) and hydroperoxyeicosatetraenoates

Target Details

Expiry Date:

12 months

rarget Details	
	(PubMed:21068195). {ECO:0000269 PubMed:12711469, ECO:0000269 PubMed:21068195}.
Molecular Weight:	55.8 kDa
UniProt:	Q96SQ9
Application Details	
Application Notes:	We expect the protein to work for functional studies. As the protein has not been tested for
	functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.