

Datasheet for ABIN7548709 **ECAT1 Protein (AA 1-217) (His tag)**



Overview

Quantity:	1 mg
Target:	ECAT1 (KHDC3L)
Protein Characteristics:	AA 1-217
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ECAT1 protein is labelled with His tag.

Product Details

Purpose:	Custom-made recombinant KHDC3L Protein expressed in mammalian cells.
Sequence:	MDAPRRFPTL VQLMQPKAMP VEVLGHLPKR FSWFHSEFLK NPKVVRLEVW LVEKIFGRGG
	ERIPHVQGMS QILIHVNRLD PNGEAEILVF GRPSYQEDTI KMIMNLADYH RQLQAKGSGK
	ALAQDVATQK AETQRSSIEV REAGTQRSVE VREAGTQRSV EVQEVGTQGS PVEVQEAGTQ
	QSLQAANKSG TQRSPEAASK AVTQRFREDA RDPVTRL Sequence without tag. The proposed
	QSEQAANNOO TONS EAASN AVTON NEDATION VITLE Sequence without tag. The proposed
	Purification-Tag is based on experiences with the expression system, a different complexit
	Purification-Tag is based on experiences with the expression system, a different complexit
Specificity:	Purification-Tag is based on experiences with the expression system, a different complexit of the protein could make another tag necessary. In case you have a special request, please
Specificity:	Purification-Tag is based on experiences with the expression system, a different complexit of the protein could make another tag necessary. In case you have a special request, please contact us.
Specificity: Characteristics:	Purification-Tag is based on experiences with the expression system, a different complexit of the protein could make another tag necessary. In case you have a special request, please contact us. If you are looking for a specific domain and are interested in a partial protein or a different

- · Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)

Grade:

custom-made

Target Details

Target:

ECAT1 (KHDC3L)

Alternative Name:

KHDC3L (KHDC3L Products)

Background:

KH domain-containing protein 3 (ES cell-associated transcript 1 protein) (KHDC3-like protein),FUNCTION: As part of the OOEP-KHDC3 scaffold, recruits BLM and TRIM25 to DNA replication forks, thereby promoting the ubiquitination of BLM by TRIM25, enhancing BLM retainment at replication forks and therefore promoting stalled replication fork restart (By similarity). Regulates homologous recombination-mediated DNA repair via recruitment of RAD51 to sites of DNA double-strand breaks, and sustainment of PARP1 activity, which in turn modulates downstream ATM or ATR activation (PubMed:31609975). Activation of ATM or ATR in response to DNA double-strand breaks may be cell-type specific (By similarity). Its role in DNA double-strand break repair is independent of its role in restarting stalled replication forks (By similarity). As a member of the subcortical maternal complex (SCMC), plays an essential role for zygotes to progress beyond the first embryonic cell divisions via regulation of actin dynamics (By similarity). Required for maintenance of euploidy during cleavage-stage embryogenesis (By similarity). Required for the formation of F-actin cytoplasmic lattices in oocytes which in turn are responsible for symmetric division of zygotes via the regulation of mitotic spindle formation and positioning (By similarity). Ensures proper spindle assembly by

Target Details

regulating the localization of AURKA via RHOA signaling and of PLK1 via a RHOA-independent process (By similarity). Required for the localization of MAD2L1 to kinetochores to enable spindle assembly checkpoint function (By similarity). Promotes neural stem cell neurogenesis and neuronal differentiation in the hippocampus (By similarity). May regulate normal development of learning, memory and anxiety (By similarity). Capable of binding RNA (By similarity). {ECO:0000250|UniProtKB:F6SZT2, ECO:0000250|UniProtKB:Q9CWU5, ECO:0000269|PubMed:31609975}.

Molecular Weight:

24.3 kDa

UniProt:

Q587J8

Application Details

Application Notes:

We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

12 months