

Datasheet for ABIN7550567

PM20D2 Protein (AA 1-436) (His tag)



Overview

Quantity:	1 mg
Target:	PM20D2
Protein Characteristics:	AA 1-436
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PM20D2 protein is labelled with His tag.

Purpose:	Custom-made recombinant PM20D2 Protein expressed in mammalian cells.
Sequence:	MRPGGERPVE GGACNGRSEL ELLKLRSAEC IDEAAERLGA LSRAIWSQPE LAYEEHHAHR
	VLTHFFEREP PAASWAVQPH YQLPTAFRAE WEPPEARAPS ATPRPLHLGF LCEYDALPGI
	GHACGHNLIA EVGAAAALGV RGALEGLPRP PPPVKVVVLG TPAEEDGGGK IDLIEAGAFT
	NLDVVFMAHP SQENAAYLPD MAEHDVTVKY YGKASHSASY PWEGLNALDA AVLAYNNLSV
	FRQQMKPTWR VHGIIKNGGV KPNIIPSYSE LIYYFRAPSM KELQVLTKKA EDCFRAAALA
	SGCTVEIKGG AHDYYNVLPN KSLWKAYMEN GRKLGIEFIS EDTMLNGPSG STDFGNVSFV
	VPGIHPYFHI GSNALNHTEQ YTEAAGSQEA QFYTLRTAKA LAMTALDVIF KPELLEGIRE
	DFKLKLQEEQ FVNAVE Sequence without tag. The proposed Purification-Tag is based on
	experiences with the expression system, a different complexity of the protein could make
	another tag necessary. In case you have a special request, please contact us.
Specificity:	If you are looking for a specific domain and are interested in a partial protein or a different
	isoform, please contact us regarding an individual offer.

Product Details

Characteristics:	Key Benefits:
	 Made to order protein - from design to production - by highly experienced protein experts. Protein expressed in mammalian cells and purified in one-step affinity chromatography The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins. State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made-to-order protein and will be made for the first time for your order. Our
	experts in the lab try to ensure that you receive soluble protein.
	If you are not interested in a full length protein, please contact us for individual protein fragments.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.
Purity:	> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)
Grade:	custom-made
Target Details	
rarget Details	
Target:	PM20D2
	PM20D2 PM20D2 (PM20D2 Products)
Target:	
Target: Alternative Name:	PM20D2 (PM20D2 Products) Xaa-Arg dipeptidase (EC 3.4.13.4) (Beta-Ala-Lys dipeptidase),FUNCTION: Catalyzes the peptide bond hydrolysis in dipeptides having basic amino acids lysine, ornithine or arginine at C-terminus. Postulated to function in a metabolite repair mechanism by eliminating alternate
Target: Alternative Name: Background:	PM20D2 (PM20D2 Products) Xaa-Arg dipeptidase (EC 3.4.13.4) (Beta-Ala-Lys dipeptidase),FUNCTION: Catalyzes the peptide bond hydrolysis in dipeptides having basic amino acids lysine, ornithine or arginine at C-terminus. Postulated to function in a metabolite repair mechanism by eliminating alternate dipeptide by-products formed during carnosine synthesis. {EC0:0000269 PubMed:24891507}.
Target: Alternative Name: Background: Molecular Weight:	PM20D2 (PM20D2 Products) Xaa-Arg dipeptidase (EC 3.4.13.4) (Beta-Ala-Lys dipeptidase),FUNCTION: Catalyzes the peptide bond hydrolysis in dipeptides having basic amino acids lysine, ornithine or arginine at C-terminus. Postulated to function in a metabolite repair mechanism by eliminating alternate dipeptide by-products formed during carnosine synthesis. {ECO:0000269 PubMed:24891507}. 47.8 kDa
Target: Alternative Name: Background: Molecular Weight: UniProt:	PM20D2 (PM20D2 Products) Xaa-Arg dipeptidase (EC 3.4.13.4) (Beta-Ala-Lys dipeptidase),FUNCTION: Catalyzes the peptide bond hydrolysis in dipeptides having basic amino acids lysine, ornithine or arginine at C-terminus. Postulated to function in a metabolite repair mechanism by eliminating alternate dipeptide by-products formed during carnosine synthesis. {ECO:0000269 PubMed:24891507}. 47.8 kDa
Target: Alternative Name: Background: Molecular Weight: UniProt: Application Details	PM20D2 (PM20D2 Products) Xaa-Arg dipeptidase (EC 3.4.13.4) (Beta-Ala-Lys dipeptidase),FUNCTION: Catalyzes the peptide bond hydrolysis in dipeptides having basic amino acids lysine, ornithine or arginine at C-terminus. Postulated to function in a metabolite repair mechanism by eliminating alternate dipeptide by-products formed during carnosine synthesis. {ECO:0000269 PubMed:24891507}. 47.8 kDa Q8IYS1

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months